Response-adapted therapy in Hodgkin's lymphoma Long-term follow up of a phase II clinical trial

Prasanth Ganesan¹, Tenali Gnana Sagar¹, Venkatraman Radhakrishnan¹, Manikandan Dhanushkodi¹, Krishnarathinam Kannan¹, Trivadi S Ganesan¹, Kadur M Lakshmipathy²
Department of Medical Oncology¹, Department of PET-CT², Cancer Institute (WIA), Chennai, India

Abstract

CONTEXT: Interim PET-CT guided therapy has emerged as one standard approaches for Hodgkin's Lymphoma (HL). We had conducted a phase II study and reported early results with 2 year follow up(Ann Oncol. 2015 Jun;26(6):1170-4) and now we present our results with more mature 5-year follow up.

DESIGN: Phase II, single-arm, single center study conducted between 2011 and 2013. Trial registration: CTRI/2012/06/002741(<http://www.ctri.nic.in>).

SETTING: Tertiary cancer center in South India.

PATIENTS: Newly-diagnosed patients with aHL.

INTERVENTIONS: Patients received 2 cycles ABVD then PET-2; if negative (Deauville score, DS 1-3) they continued 4 additional cycles of ABVD and if positive (DS 4, 5) they received 4 cycles of Escalated BEACOPP

MAIN OUTCOMES MEASURES: Primary end point was event free survival (EFS) – from start of therapy till event (relapse after initial response, or progression from PR or death). The secondary end points were overall survival and toxicity of EB.

RESULTS: Fifty patients [median age: 28 years, 12-60; 78% males; Stages: IIB: 3 (6%), III: 29 (58%) and IV: 18 (36%); High risk IPS score 4-7: 16 (32%)] were included and 49 had interim PET study [positive 8 (16%), negative 41]. EB was started in 7/8 patients with positive scans, 5 (of 7) achieved complete remission 4 remain event-free till now. Among PET-2-ve patients (N=41), 34 remain event free. After a median follow up of 66 months (3-89), the estimated 5-yr EFS and OS was 77% and 85% respectively for the entire cohort of 50 patients. For the PET-2 positive patients, 5-yr EFS and OS was 50% and 62% and for the PET-2 negative group it was 82% and 89% respectively.

CONCLUSIONS: The results of response-adapted escalation treatment strategy in aHL are sustained and the outcomes of PET-2 positive patients are improved. Outcomes in PET-2 negative patients are lower than those reported in retrospective analyses but comparable to similar prospective studies.

Background

- 5 yr-EFS in Advanced Hodgkin's Lymphoma (stage IIB, III, IV)
 - ABVD: 61-73%^{1,2}
 - Esc BEACOPP (EB): 85%³
- EB is more toxic than ABVD³
 - Febrile neutropenia 22% vs. 0.5%
 - AML/MDS at 10 years: 3.2% vs. <1%
- Interim PET-CT after 2 cycles of ABVD (PET-2)^{4,5} is a robust predictor of survival in advanced HL
 - PET-2 positive EFS: 12-28%
 - PET-2 negative EFS: 94-95%
- PET- guided strategy can selectively tailor high intensity therapy for the highest risk patients

Materials & Methods

RESEARCH QUESTION

- Can interim-PET (after 2 cycles of ABVD, PET-2) guided intensification of therapy (PET-2 positive patients to escalated BEACOPP) improve outcomes in advanced (stages IIB, III, IV) HL?

DESIGN

- Phase II, single-arm, single center study conducted between 2011 and 2013
- Fleming's one-stage design- sample size 50 patients**
 - Type I error $\alpha=0.05$ and power= 0.8
 - p0: 70%, p1: 85%

INCLUSION

- Newly diagnosed advanced HL (Stage IIB, III, IV)
- Age 12-65 years
- Normal organ function

Baseline characteristic	N	%
Age, years		
Median	28	
Range	12-60	
Male sex	39	78
ECOG PS		
0-1	47	94
2-3	3	6
Stage		
IIB	3	6
III	29	58
IV	18	36
B symptoms present	32	64
Bulky disease present	12	24
IPS score		
0-3	34	68
4-7	16	32

END POINTS

- Event free survival (EFS) – from start of therapy till event (relapse after initial response, or progression from PR or death).
- Overall survival (OS)- from start of therapy till death due to any cause

Results

Parameter	N	5-year EFS%	P value
B symptoms			
Absent	18	69	0.34
Present	32	81	
Bulky disease			
Absent	38	73	0.26
Present	12	91	
IPSS			
0-3	34	82	0.21
4-7	16	67	
Interim PET			
Negative	41	82	0.019
Positive	8	50	

Entire cohort of 50 patients (intention to treat analysis)

5-yr EFS 77%

5-yr OS 85%

- Of the 3 patients who failed EB, none could be salvaged
- Among the 5 who failed ABVD, all 5 responded to salvage chemotherapy, and 4 underwent autologous transplant
- There are no cases of secondary cancers till now

Discussion

- iPET 2 positive patients improved their survival with escalation of therapy when compared to historical data
- PET-2 guided strategy is able to deliver tailored therapy and avoids toxicity of EB in majority of the patients

References

- Canelos P. J Clin Oncol. 2014;32(3):163-8
- Ganesan P. Ann Hematol. 2011;90(10):1153-60
- Diehl V. N Engl J Med. 2003;348(24):2386-95
- Gallamini A. J Clin Oncol. 2007;25(24):3748-52
- Gallamini A. Haematologica. 2014;99(6):1107-13
- Ganesan P. Ann Oncol. 2015 Jun;26(6):1170-4

Trial registration: CTRI/2012/06/002741(<http://www.ctri.nic.in>).