PCSK9-inhibition in routine clinical practice: real world data from Germany

E. Waldmann, J. Altenhofer, K. Henze, K.G. Parhofer

Introduction

PCSK9-inhibitors (PCSK9-i) are a new treatment option for patients not reaching their LDL-cholesterol targets. Efficacy and safety is good in study settings, but there is only few data on real world experience. We analyzed data from 146 consecutive patients who received PCSK9-i and participated in an observational study (lipid clinic; monocentric).

Methods

Lipid profiles were obtained (age 63 ± 10 y, BMI 23.6 ± 4.1 kg/m2, 49% male) before and after 3-4 months treatment with PCSK9-I.

	baseline (mg/dl)	after 3 months (mg/dl)	change (%)	p-value
cholesterol	250.3 ± 72.1	161.5 ± 51.7	-34.4 ± 14.5	<0.001
triglycerides	168.5 ± 102.8	163.0 ± 99.5	4.1 ± 42.5	0.440
HDL-cholesterol	55.9 ± 15.4	55.5 ± 15.3	-0.2 ± 11.0	0.378
Non-HDL-cholesterol	194.4 ± 70.4	106.0 ± 49.4	-44.7 ± 18.0	<0.001
LDL-cholesterol	161.0 ± 58.9	80.1 ± 42.7	-49.9 ± 18.8	<0.001
VLDL-cholesterol	33.7 ± 42.0	26.0 ± 27.4	1.9 ± 87.1	0.001
VLDL-triglycerides	145.9 ± 104.7	143.1 ± 99.7	8.7 ± 58.9	0.499
lipoprotein(a)	48.6 ± 48.5	39.8 ± 40.6	-24.2 ± 20.4	<0.001

Fig.1: Parameters of lipid metabolism before and under 3 months PCSK-9-i therapy in mg/dl

	baseline (mg/dl)	after 3 months (mg/dl)	change (%)
Alirocumab			
LDL-cholesterol	158.2± 55.6	80.6± 42.7	-48%± 19
lipoprotein(a)	52.2± 57.5	43.7± 46.6	-23%± 23
Evolocumab			
LDL-cholesterol	174.7± 80.1	89.6± 67.1	-50%± 21
lipoprotein(a)	46.8± 63.7	41.6± 58.5	-24%± 25

Fig.2: Parameters of lipid metabolism comparing Evolocumb and Alirocumab in mg/dl

Results

44% received Evolocumab 140mg and 56% Alirocumab (29% 75mg and 27% 150mg) q 2 weeks. Patients had either (95%) atherosclerotic disease or familial hypercholesterolemia without atherosclerosis (5%). All were on maximally tolerated lipid lowering therapy. Statin intolerance was prominent (complete 44%, partial 25%). Baseline LDL-c was reduced by 50% (Fig.1); there was no significant difference between Alirocumab and Evolocumab (Fig.2). 17 (12%) patients had a LDL-c reduction of <30%. In 8 lipid lowering therapy was reduced, in 4 (2.7%) no reason was detectable, 3 varied considerably with stronger reductions at earlier or later time points. In 2 injection was longer than 2 weeks ago. Treatment was tolerated well. No patient stopped PCSK9-i within the first 3 months.

Conclusions

Alirocumab and Evolocumab reduce lipids safely in routine clinical practice. Statin intolerance is one of the main reasons to initiate PCSK9-inhibitors. True non-responders are rare (<3%). Baseline LDL-C is considerably higher than in outcome studies (161 mg/dl vs. 90mg/dl). Whether this translates into even greater clinical benefit than in the studies remains to be evaluated.