Review of Practice Patterns in Post-treatment Surveillance of Primary Epithelial Ovarian Cancer

Authors: Joseph DeMari¹, MD; Monica Hagen Vetter², MD; Shruthi Chandra⁴; John Hays, MD PhD³; Ritu Salani, MD MBA²

Department of Obstetrics and Gynecology, The Ohio State University Wexner Medical Center¹; Division of Gynecologic Oncology, The Ohio State University Comprehensive Cancer Center – James Cancer Hospital and Solove Research Institute²; Division of Medical Oncology, The Ohio State University Comprehensive Cancer Center – James Cancer Hospital and Solove Research Institute³; The Ohio State University College of Medicine⁴

The James

Objective

To determine practice patterns for surveillance of primary ovarian cancer after complete response to therapy and to identify the percentage of clinicians who follow the surveillance guidelines endorsed by the Society for Gynecologic Oncology (SGO).

Methods

This is a single-institution retrospective chart review of all patients with epithelial ovarian cancer with a complete response to primary therapy between January 2012 and January 2015. Descriptive statistics were performed due to the exploratory nature of the study.

Discussion

The vast majority of patients are receiving surveillance care compliant with SGO guidelines. Adherence to the SGO guidelines ensures cost-reducing surveillance as the guidelines eliminate the need for routine CT imaging.

References

- Salani, R., Khanna, N., Frimer, M., Bristow, R. E., & Chen, L.-M. (2017). An update on post-treatment surveillance and diagnosis of recurrence in women with gynecologic malignancies: Society of Gynecologic Oncology (SGO) recommendations. *Gynecologic Oncology*, 146(1), 3–10.
- Salani, R., Backes, F. J., Fung, M. F. K., Holschneider, C. H., Parker, L. P., Bristow, R. E., & Goff, B. A. (2011). Posttreatment surveillance and diagnosis of recurrence in women with gynecologic malignancies: Society of Gynecologic Oncologists recommendations. *American Journal of Obstetrics and Gynecology*, 204(6), 466–478.
- Rimel, B. J., Burke, W. M., Higgins, R. V., Lee, P. S., Lutman, C. V., & Parker, L. (2015). Improving quality and decreasing cost in gynecologic oncology care. Society of gynecologic oncology recommendations for clinical practice. *Gynecologic Oncology*, 137(2), 280–284.

SGO Guidelines

	Years 0-2	Year 2-3	Years 3-5	Years >5
Review of symptoms and physical exam	3-4 months	4-6 months	6 months	Yearly*
Pap test/cytologic evidence	Not indicated	Not indicated	Not indicated	Not indicated
CA-125	Optional	Optional	Optional	Optional
Radiographic imaging (CXR, PET/CT, MRI, US)	Insufficient data to support routine use	Insufficient data to support routine use	Insufficient data to support routine use	Insufficient data to support routine use
Recurrence suspected	CT scans or PET/CT scans CA-125	CT scans or PET/CT scans CA-125	CT scans or PET/CT scans CA-125	CT scans or PET/CT scans CA-125

*may be followed by a gynecologist oncologist or generalist

Results

Figure 1: Shows percentage of patients that were imaged for a given indication. The majority of patients were imaged due to rising Ca-125 levels, symptoms concerning for recurrence, or both.

Figure 2: Shows method of detection of cancer recurrence. The majority of recurrence was detected via rising Ca-125 levels.

Follow-up

- 82% had clinical follow-up scheduled according to SGO guidelines (+/- 1 month)
- Mean time to recurrence 11.8 months

Compliance

- 18% had abnormal follow-up that was non-compliant with SGO guidelines
 - 3/9 were due to patient initiated visits/point of access visits
 - 6/9 were due to unknown reasons
- Stage and histology were not associated with with non-compliance or frequency of follow-up