MULTIMODAL OPIOID-SPARING ANALGESIA PROTOCOL IN COLORECTAL LAPAROTOMIC SURGERY: when epidural catheter is not possible.

Á. Herrero Cano¹, F. Muriedas Alfonso¹, C. Sánchez- Cabezudo Moreno¹, J. Casado Neira¹, M. Barbero Mielgo¹, J. García Fernández¹.

¹Anaesthesia, Critical care and Pain Department. Puerta de Hierro University Hospital. Majadahonda, Madrid.

BACKGROUND:

ERAS (Enhance Recovery After Surgery) decrease surgical damage, resulting in the reduction of morbidity, mortality and length of hospital stay. Epidural analgesia is gold standard in laparotomic surgeries, however sometimes is not possible. We analysed many data in order to find out if multimodal analgesia is effective also without epidural catheter.

MATERIALS AND METHODS:

We collected laparotomic colorectal (CCR) surgical patients without epidural catheter. Group A (N: 25) were submitted in ERAS program in 2017 and group B (N: 16): an historic group without ERAS program in 2015.

We did not place an epidural catheter due to contraindication, technical impossibility or laparoscopies converted to laparotomies.

	PREOPERATIVE	INTRAOPERATIVE	POSTOPERATIVE
GROUP A	Gabapentine 300 mg.	- Magnesium IV (30 mg/kg) - Lidocaine IV (bolus 1,5mg/kg + continuous infusion 1mg/kg). - Dexamethasone (0,1 mg/kg). - NSAID. - Bilateral transversus abdominis plane (TAP) block.	- NSAID. - Morphine 0.5 mg/kg if VAS>4 EPIDURAL CATHETER was not placed.
GROUP B		- Opioids and NSAID. - Dexamethasone (0,1 mg/kg).	

RESULTS:

VAS AFTER SURGERY (p>0.05)

MORPHINE CONSUMPTION AFTER SURGERY mg (p<0.05)

POVN – ILEUS- HOSPITAL STAY (p>0.05)

REACTIVE C PROTEIN (RCP) AFTER SURGERY (p>0.05)

CONCLUSIONS:

Multimodal opioid- sparing analgesia program seems to decrease opioids consumption, side effects, RCP and length of hospital stay in CRS patients even if epidural analgesia is not possible. In order to get significative results the sample should be extended.