Retrospective audit into the Management of Epistaxis in the Emergency department

E Tyler, S Khwaja

INTRODUCTION

Epistaxis is a common ENT condition accounting for 1 in every 200 visits to the Emergency Department (ED). It has the potential to be life-threatening if not treated appropriately and in a timely manner. Causes of epistaxis are varied, ranging from nasal trauma to neoplasia in rare cases, however, it most commonly occurs spontaneously without prior insult. Those particularly at risk include those with hypertension and the anticoagulated elderly population. Management of epistaxis consists of application of nasal pressure to encourage haemostasis naturally, cautery of bleeding points with topical silver nitrate and nasal packing for refractory cases, surgical intervention is rarely indicated.

METHODS

A retrospective audit was conducted within the ED at Stepping Hill Hospital over a 4 week period, to identify all adults presenting with epistaxis and measure the management modalities. These were analysed against

predetermined criteria to assess the appropriateness and efficacy of management. Further in-depth review was conducted of 6 cases where the indications for nasal packing were unclear.

Management Modalities

RESULTS

In total, 26 patients were included, nasal pressure was applied in 75% of patients who were actively bleeding on presentation, 41.7% were cauterised where it was indicated and 62.5% of those packed were appropriate and 37.5% being deemed as inappropriately packed, no patients that did not receive packing required it.

Audit Standards

	Criterion	Results		Meets/Exceeds Standard
		Number	Percentage	
1	Nasal pressure applied	15/20	75.0%	No
2	Cautery attempted	5/12	41.7%	No
3	Packing attempted	5 Appropriate, 3 Potentially inappropriate	Not measurable	N/A
4	Blood tests performed	17/18	94.4%	No
5	Sufficient Documentation	15/26	57.7%	No

Bleeding Duration whilst in hospital

CONCLUSION

Recommendations were made to increase the utilisation of conservative methods of managing epistaxis such as cautery and decongestant spray, in order to reduce the rates of nasal packing and thus admission to hospital and overnight stays.

1. Stocking of phenylcaine spray within the ED given by Triage Nurses
2. Training of Nurse Practitioners in silver nitrate cautery
3. Development of a local epistaxis management protocol (figure below)