Hyperhomocysteinemia is not associated with systemic arterial stiffness in patients with acute ischemic stroke

Kwang-Yeol Park, MD, PhD; Jeong-Min Kim, MD, PhD

Department of Neurology, Chung-Ang University Hospital, Chung-Ang University College of Medicine, Seoul, Korea

Background:

Hyperhomocysteinemia is reported to be a risk factor for vascular calcification, atherosclerosis, and cerebrovascular disease. Also, arterial stiffness is associated with increased pulse pressure and cerebral small vessel disease. The aim of this study was to determine the association between hyperhomocysteinemia and brachial-ankle pulse wave velocity (baPWV) in patients with stroke.

Methods:

We identified consecutive patients with acute ischemic stroke who visited a tertiary university hospital within seven days of symptom onset between January 1, 2011 and April 30, 2017. The hyperhomocysteinemia was defined as serum level of homocysteine > 11 umol/L. The average of baPWV measured on both sides was categorized into quartiles. The association between homocysteine and baPWV was tested using chi-square test and logistic regression analysis.

Results:

A total of 1624 patient were enrolled in this study. Mean age was 68.5 +/- 12.5 years and male was 882 (54.3%). Hyperhomocysteinemia was found in 1074 (66.1%). On bivariate analysis, baPWV was significantly associated with age, sex, hypertension, diabetes mellitus, smoking, systolic blood pressure, highly sensitive C reactive protein, estimated GFR, and hyperhomocysteinemia. However, the association between hyperhomocysteinemia and baPWV was not significant on multivariable logistic regression. The interaction term of hyperhomocysteinemia and sex was not significant, either. Age, hypertension, diabetes mellitus, and systolic blood pressure remained significant on multivariable analysis.

Conclusion:

Despite previous studies showed the association between homocysteine and arterial stiffness in general population, our study did not show significant association between them in stroke population. We presumed that potent vascular risk factors such as hypertension and diabetes rather than homocysteine play a more important role in damaging vessel wall in stroke patients. Further studies are warranted to confirm our findings and hypothesis.

Table 1. Baseline information

	Total	Q1	Q2	Q3	Q4	p_value
Age	68.46 +/- 12.46	59.06 +/- 12.81	67.09 +/- 10.96	71.23 +/- 9.77	76.48 +/- 8.92	0.000
Sex	882 (54.3%)	268 (66%)	249 (61.2%)	209 (51.6%)	156 (38.4%)	0.000
HTN	969 (59.7%)	150 (36.9%)	247 (60.7%)	265 (65.4%)	307 (75.6%)	0.000
DM	482 (29.7%)	65 (16%)	120 (29.5%)	138 (34.1%)	159 (39.2%)	0.000
IHD	160 (9.9%)	37 (9.1%)	46 (11.3%)	43 (10.6%)	34 (8.4%)	0.479
Af	341 (21%)	74 (18.2%)	78 (19.2%)	96 (23.7%)	93 (22.9%)	0.146
Smoking	414 (25.5%)	150 (36.9%)	129 (31.7%)	95 (23.5%)	40 (9.9%)	0.000
HL	214 (13.2%)	49 (12.1%)	50 (12.3%)	63 (15.6%)	52 (12.8%)	0.428
PrevCVA	272 (16.7%)	53 (13.1%)	69 (17%)	75 (18.5%)	75 (18.5%)	0.124
NIHSS	4.56 +/- 5.73	4.3 +/- 5.89	4.45 +/- 5.78	4.41 +/- 5.43	5.09 +/- 5.81	0.203
SBP	151.25 +/- 27.69	142.54 +/- 25.39	150.96 +/- 25.1	153.43 +/- 27.03	158.06 +/- 30.66	0.000
DBP	86.74 +/- 14.22	84.87 +/- 13.95	86.8 +/- 13.44	86.87 +/- 14	88.45 +/- 15.26	0.005
hsCRP	7.26 +/- 21.28	5.15 +/- 19.19	5.9 +/- 15.28	7.36 +/- 20.62	10.62 +/- 27.76	0.001
FBS	135.29 +/- 56	126.58 +/- 54.12	133.75 +/- 49.19	139.68 +/- 58.35	141.17 +/- 60.72	0.001
HbA1c	6.21 +/- 1.38	5.92 +/- 1.25	6.23 +/- 1.47	6.37 +/- 1.44	6.33 +/- 1.31	0.000
BUN	18.13 +/- 8.22	16.44 +/- 6.03	18.09 +/- 7.9	18.57 +/- 8.4	19.43 +/- 9.83	0.000
Cr	0.98 +/- 0.82	0.9 +/- 0.6	0.96 +/- 0.64	1.06 +/- 1.17	0.99 +/- 0.77	0.043
eGFR	87.38 +/- 27.34	102.26 +/- 24.3	88.7 +/- 26.78	82.19 +/- 25.32	76.34 +/- 25.97	0.000
T_chol	179.34 +/- 45.09	180.3 +/- 43.13	179.02 +/- 42.79	178.19 +/- 45.61	179.85 +/- 48.69	0.915
HDL	46.36 +/- 13.29	46 +/- 11.29	45.68 +/- 13.47	46.78 +/- 12.95	47 +/- 15.17	0.446
LDL	106.35 +/- 33.23	107.27 +/- 31.96	108.7 +/- 32.11	104.39 +/- 33.71	105.03 +/- 34.98	0.223
TG	117.54 +/- 77.05	119.56 +/- 90.64	116.36 +/- 68.74	116.47 +/- 72.63	117.76 +/- 74.61	0.928
SVO	331 (22.9%)	74 (22.2%)	78 (21.5%)	92 (24.9%)	87 (22.8%)	0.718
Homocystein	14.01 +/- 6.95	13.62 +/- 7.11	14.23 +/- 8.43	14.31 +/- 6.48	13.9 +/- 5.43	0.467
Homocystein_High	1074 (66.1%)	254 (62.6%)	255 (62.7%)	278 (68.6%)	287 (70.7%)	0.024
VitD	14.84 +/- 8.48	15.5 +/- 8.46	14.79 +/- 7.6	14.8 +/- 8.02	14.26 +/- 9.69	0.258
VitD_adj	14.82 +/- 8.33	15.52 +/- 8.3	14.83 +/- 7.32	14.7 +/- 8.01	14.23 +/- 9.52	0.207

Table 2. Multivariable analysis

	Estimate	Std. Error	t value	Pr(> t)	2.5 %	97.5 %
(Intercept)	-102.8261377	144.1666924	-0.7132448	0.4758074	-385.6203473	179.968072
Age	21.0968699	1.2266007	17.1994607	0.0000000	18.6907970	23.502943
SexM	-76.3997440	27.6731318	-2.7607914	0.0058378	-130.6827500	-22.116738
eGFR	0.2550936	0.5807549	0.4392449	0.6605486	-0.8841024	1.394290
HTNHT(+)	130.7906977	26.6190895	4.9134174	0.0000010	78.5752779	183.006117
DMDM(+)	115.2098656	27.5637650	4.1797579	0.0000309	61.1413911	169.278340
SmokingSmoking(+)	-54.0549188	31.6355737	-1.7086752	0.0877221	-116.1105637	8.000726
SBP	4.4923627	0.4410054	10.1866378	0.0000000	3.6272961	5.357429
hsCRP	1.4951541	0.5834227	2.5627286	0.0104841	0.3507249	2.639583
VitD_adj_3gr1	-71.3896158	28.6658476	-2.4904066	0.0128693	-127.6199113	-15.159320
VitD_adj_3gr2	-26.8295846	35.1092082	-0.7641752	0.4448854	-95.6990343	42.039865
Homocystein_High	13.1124857	28.7318694	0.4563743	0.6481882	-43.2473166	69.472288