Taxane and platinum chemotherapy for patients with uterine carcinosarcoma

Hiroyuki Yamazaki, Yukiharu Todo, Chisa Shimada, Shinichiro Minobe, Hidenori Kato
Division of Gynecologic Oncology, National Hospital Organization, Hokkaido Cancer Center, 4-2 Kikusui, Shiroishi-ku, Sapporo 003-0804, Japan

Objectives:

To assess the effects of taxane and platinum combined chemotherapy in uterine carcinosarcomas(UCS).

Methods:

We retrospectively examined the records of women with UCS and overall survival(OS)/survival after recurrence(SAR) are demonstrated in Kaplan-Meier method. And disease control rates of chemotherapies were evaluated in women who had measurable disease.

Table. Characteristics.

*Including adjuvant chemotherapies without measurable diseases

	All eligible patients	With measurable diseases
No. of patients	45	28
Advanced	29	19
Recurrence	16	9
Age		
median	64	62
range	47-82	47-77
Stage		
IA	9	4
IB	3	2
II	4	3
IIIA	2	0
IIIC1	4	1
IIIC2	4	2
IVB	19	16
Surgery		
Biopsy only	3	3
Hysterectomy	18	13
+LND/LN biopsy	24	12
pN		
Nx	21	16
N0	15	9
N1	9	3
Previous Taxan+Platinum*		
Yes	25	21
No	20	7

Results:

45 women were eligible among seventy-five patients with UCS in 1987-2016. 29 had advanced stage (III or IV) and 16 had recurrence disease. Their overall survival and survival after recurrence was 8 months (median, range 1-45) and 10 months (median, range 1-68). The two-year survivals were 20% and 22%, respectively. 28 patients had measurable disease when treated with chemotherapy, which accounted for 39 regimens. The disease control rate of taxane and platinum combined chemotherapy was 38.9%.

Conclusions:

Taxane and platinum combined chemotherapy for advanced or recurrent UCS showed less activity than previously reported although this is a single center retrospective study in the small number.

Figure. Number of the patients.

Figure. Kaplan-Meier survival curves.

:OS in advanced group, SAR in recurrence group.

Table. Response to the chemotherapies grouped by the key-drugs

	N=	CR	PR	SD	PD	Previous chemotherapy
		CR+PR+SD (Disease control rate)				
Taxane and platinum	18	2	2	3	11	3 (16.7%)
		7 (38.9%)				
Gemcitabine	8		1		7	8 (100%)
		1 (12.5%)				
Adriamycin	6			1	5	4 (66.7%)
		1 (16.7%)				
Platinum (without taxane)	4		1		3	2 (50%)
		1 (25%)				
Pazopanib	3		1	1	1	2 (66.7%)
		2 (66.7%)				