MANAGING THERAPEUTIC RELATIONSHIP IMPASSES IN CBT

D. Kljenak, Department of Psychiatry, University of Toronto

Introduction

Cognitive behaviour therapy (CBT) is an evidence-based therapy used to effectively treat a number of mental health disorders. Traditionally, CBT has been thought of as a treatment modality that is technique-based and not as concerned with the therapeutic relationship than other forms of psychotherapy. Actually, CBT recognizes that a good therapeutic relationship is essential for effectively collaborating with any patient and especially with patients with complex diagnostic presentations, such as patients with personality disorders. The second Task Force on Evidence-Based Therapy Relationships concludes that “The therapy relationship makes substantial and consistent contributions to psychotherapy outcome independent of the specific type of treatment” and that “The therapy relationship accounts for why clients improve (or fail to improve) at least as much as the particular treatment method.” It has also been recognized that optimal handling of therapeutic alliance ruptures are cornerstone of any effective therapy including CBT.

In order to effectively address potential alliance ruptures therapist needs to be aware of interpersonal processes (transference and countertransference). Teaching the trainee CBT therapist strategies to address countertransference is important in preventing poor outcomes in supervision and therapy.

Teaching CBT Strategies to Address Countertransference

1. Acknowledge countertransference exists and is a valuable source of information about the patient’s current and past relationships.
2. Develop self-awareness (awareness of personal schemas and underlying assumptions and awareness of personal triggers for your own schema activation).

- Practice CBT on oneself
- Supervision/consultation
- Undergo own therapy

A. Practicing CBT on oneself:

- Monitor (therapist’s) own subtle physical sensations (i.e. muscle tensing), mood shifting (i.e. anxiety about interrupting the patient) and behaviour, especially being alert for different-from-the-typical behaviour (i.e. while able to set limits with other patients, having difficulty setting limits with one particular patient)
- Use above as a reminder that personal schema may be activated
- Complete a dysfunctional thought record (in other words, identify one’s own automatic thoughts and develop a rational response
- Identify one’s own underlying beliefs (UA) about the patient, self and the therapeutic relationship
 - By using standard (UA) identification methods
 - By completing the Dysfunctional Attitude Scale (Weissman & Beck, 1978)
- Identify one’s own core beliefs/schemas
 - By using standard schema identification methods such as downward arrow technique, looking for central themes in own automatic thoughts, directly eliciting schemas about self, the patient and the therapeutic relationship
 - By completing the Therapist’s Schema Questionnaire (Leahy, 2001), the Personal Belief Questionnaire (Beck and Beck, 1995), and the Young Schema Questionnaire (Young & Brown, 2001)

B. Asking the following questions:

- Is my response due to my personal schema activation by the patient’s behaviour?
- Am I entangled in the “self-fulfilling prophecy” (Leahy, 2009) game initiated by the patient?
- Does this fit with what I learned about the patient’s developmental history/early relationships with significant others (early attachment relationships) (i.e. am I reacting the way the patient’s early attachment figures were reacting or am I reacting the way the patient was responding to their early attachment figures)?
- Does this fit with the patient’s current interpersonal relationship (i.e. if I am feeling this way/wanting to behave this way – do other people in the patient’s life react in a similar way)?
- What is the patient’s response to this significant other reaction or, in other words, what is the function of this behaviour for the patient?

Case Example

Dr. A. realizes that the thought of following an agenda and ending sessions on time with Mr. B. makes her anxious. She decides to explore her beliefs about therapy/self/patient (Rudd and James, 1997).

Belief about the Patient

- The patient will not like structured therapy.
- The patient will not like me if I attempt to structure therapy

Belief about Self

- I should be liked by the patient
- I should please the patient
- I should meet my patient needs

Beliefs about the course of Therapy

- CBT will not work with this patient

Therapist’s behavioural response

- Allows the patient to ramble, does not set or follow agenda; extends sessions over time

Dr. A. realizes that she is:

- Using a number of cognitive distractions i.e. fortune-telling, should statements, personalizing, over-generalization
- The above beliefs represent her own need for approval and excessive self-sacrifice schema activation that leads to unatypical-for-her behaviour in the session.

Case Example cont.

- . In addition, she realizes that she is mirroring the patient’s behaviour (the patient has difficulty setting limits at her work and repeatedly works unpaid overtime).
- Dr. A. asks herself about advantages/disadvantages of agenda setting with this patient.
- Dr. A. decides to do a behavioural experiment in her next session with the patient to test her assumption that the patient will not like structured therapy.
- At the next session, the patient gives her very positive feedback about the session (“I liked this session. It makes me feel we are going somewhere. I liked that we set an agenda”)

Therapist’ s Schemas - Reflection :

- Are there some patients that induce strong feelings in you?
- What beliefs, interpersonal schemas, and behaviours do you think have been activated?
- Potential difficulties with therapist and patient schemas match/mismatch?

Alliance rupture

Cognitive Interpersonal Cycle

- Hypothesis about the patients interpersonal beliefs, expectations, and appraisal processes that contribute to the perpetuation of unhelpful cognitive-interpersonal cycles (Safran & Segal, 1990).
- Cycle maintained by cognitive schema

- Therapeutic intervention activates an important IP schema ALLIANCE RUPTURE
- Opportunity to explore and modify IP schema

Alliance repair

- Importance of therapist’ s reactions and interpersonal schemas as therapy tools
- Distinguish between being hooked and unhooked (to be hooked means not to be aware of one’s feelings and IP tendencies and thus unwittingly participate in the interaction)
 - Are your responses complementary or not to the patient’s interpersonal pool
 - Examine own feelings, action tendencies and to which interpersonal markers you respond

- Applying DECENTERING - Unhooking from Pt’s cognitive - interpersonal cycle
- Non-judgmental Awareness & Attention to:
 - Own thoughts / feelings
 - Patient’ s thoughts / feelings
 - Therapeutic process (vs content)
- EXPERIENTIAL DISCONFIRMATION Use the therapeutic relationship to disconfirm dysfunctional beliefs
 - Collaborative exploration, empathy
 - Meta-communication
- Identification and Exploration of:
 - Therapist’s / patient’s – thoughts / feelings
 - Links to the here-and-now processes and relationship
 - Links to the hypothesized cognitive-interpersonal cycle
 - Links to the past
 - Links to behaviors outside the session
- Therapist -> owning up to own contribution
- May involve therapist self-disclosure

Summary

Cognitive behaviour therapy (CBT) is an evidence-based therapy used to effectively treat a number of mental health disorders. Traditionally, CBT has been thought of as a treatment modality that is technique-based and not as concerned with the therapeutic relationship than other forms of psychotherapy. Increasingly it has been recognized that a good therapeutic alliance and optimal handling of therapeutic alliance ruptures are cornerstone of any effective therapy including CBT. Transference and countertransference need to be addressed in order to effectively prevent and deal with therapeutic impasses which can be understood through a CBT lens. Cognitive and behavioural techniques could be effectively used to recognize and manage challenging moments in therapeutic relationship. This is important in preventing poor outcomes in therapy.

References

1. Beck, A., & Beck, J. (1995). The personality belief questionnaire. Bala Cynwyd, PA: Beck Institute for Cognitive Therapy and Research.
2. Bennett-Levy, J., Turner, F., Beatty, T., Smith, M., Paterson, B., & Farmer, S. (2001). The value of self-practice of cognitive therapy techniques and self-reflection in the training of cognitive therapists. *Behavioural and Cognitive Psychotherapy*, 29, 203-220.
3. Bernard, J.M., Goodyear, R.K. (1992). Fundamentals of clinical supervision, Weecham Heights, MA: Allyn & Bacon
4. Gluhoski, V.L., (1994). Misconceptions of cognitive therapy. *Psychotherapy*, 31, 594-600.
5. Haarhoff, B. (2006). The importance of identifying and understanding therapist schema in cognitive therapy training and supervision. *New Zealand Journal of Psychology*, 35(3), 126-131.
6. Liese, S.B., Barber, J.P., Beck A.T. (1995.) The Cognitive Therapy Adherence and Competence Scale. Unpublished instrument, University of Kansas Medical Center.
7. Liese, S.B., & Beck, J.S.(1997). Cognitive therapy supervision. In C.W. Watkins (Ed.), *Handbook of psychotherapy supervision* (pp. 114–133). New York: Wiley
8. Leahy, R.L. (2001). *Overcoming resistance in cognitive therapy*. New York: The Guilford Press.
9. Leahy, R.L. (2009). Schematic mismatch in the therapeutic relationship. A social-cognitive model. In P.Gilbert & R.L. Leahy(Ed.) The therapeutic relationship in the cognitive behavioral psychotherapies (pp.229-254). New York: Routledge
10. Rudd, M., & Joiner, T. (1997). Countertransference and the therapeutic relationship: A cognitive perspective. *Journal of Cognitive Psychotherapy: An International Quarterly*, 11(4), 231-249.
11. Weissman, A., & Beck, A. (1978, March 27-31). *Development and validation of the Dysfunctional Attitude Scale: A preliminary investigation*. Paper presented at the Annual Meeting of the American Educational Research Association, Toronto, Canada
12. Young, J.E., & Brown, G. (2001). *Young schema questionnaire*. New York: Cognitive Therapy Center.
13. Safran JD, Segal ZV. Interpersonal process in cognitive therapy. New York: Basic Books; 1990.