CLOSTRIDIUM DIFFICILE INFECTION IN CHILDREN: EXPERIENCE IN A TERTIARY CARE HOSPITAL

A. Calvo Cillán¹, L. Martin Pedraz¹, B. Carazo Gallego², D. Lopez Martin¹, C. Sole Mesalles³, B. Palop Borrás⁴, D. Moreno Perez².

¹Hospital Materno Infantil de Malaga, Pediatrics, Málaga, Spain. ²Hospital Materno Infantil de Malaga, Pediatric Infectology and Immunodeficiencies Unit, Málaga, Spain.

³Complexo Hospitalario Universitario de Ourense, Pediatrics, Ourense, Spain. ⁴Regional University Hospital of Málaga, Microbiology, Málaga, Spain.

Background

Clostridium difficile infection (CDI) has increased in children, although its diagnosis in <3 years-old is controversial due to high rates of colonization. Retrospective study to describe the characteristics of CDI diagnosed in children <16 years-old during 2013-2017. CDI in <1 years-old were excluded following the recommendations of the literature.

Case Presentation Summary

Over the period, 30 episodes of CDI were diagnosed. 2 cases were excluded for being <1 years-old. The remaining 28 were diagnosed in 23 children. Mean age at diagnosis: 6.79 years (SD 4.68). 39.1 % were between 1-3 years-old. Risk factors (RF) for CDI are presented in *figure 1*. Abdominal pain (57.9%), diarrhea (46.3%) and fever (10.5%) were the predominant symptoms. 75% had mild-moderate colitis, no one presented as fulminant colitis. CD toxin was positive in 67.9% and Nucleic-Acid Amplification Test in 78.6%; both resulted positive in 46.4%. Most of the first episodes were treated with metronidazole (68.4%); rifaximine was used in 14.3% of them. The length of treatment was 10 days in 68 %. Two patients were not treated (7.1%), with good outcome. Six children developed recurrences (22.2%), generally treated with metronidazole (33%). One patient presented multiples recurrences and fidaxomicin therapy was indicated, without more relapses. The use of gastric acid-suppressing agents (GASA) or a previous episode of CDI was associated with the development of recurrence ($p<0.05$ and $p=0.06$). No statistically association were observed between other RF and recurrence.

Learning Points

- Most CDI presented with mild-moderate colitis, as in other paediatric studies.
- 39.1% CDI happened in <3 year-old children, although some of them could be simple colonization.
- Use of Gastric acid-suppressing agents was associated with relapsing CDI.

Risk Factors of CDI (n=28)	
RF	n (%)
Immunosuppressive treatment	14 (50%)
Previous hospitalization	9 (32,1%)
Inflammatory bowel disease	8 (28,6%)
Gastric acid-suppressing agents	8 (28,6%)
Previous antibiotherapy	8 (28,6%)
Previous CDI episode	6 (21,4%)
Broad spectrum antibiotic therapy	6 (21,4%)
Oncologic diseases	5 (17,9%)

Figure 1. Risk factors of CDI. RF: Risk Factors; CDI: *Clostridium difficile* Infection