Ultrasound-guided Selective Versus Conventional block of the Medial Brachial Cutaneous and the Intercostobrachial Nerves.

P. Magazzeni, H. Bouaziz, G. Mekler, E. Albuissou, D. Jochum, G. Iohom

Background and Goal of Study :

- Anesthesia for superficial surgery of antero- and posteromedial surfaces of the upper arm :
 - Axillary brachial plexus block (ABPB) AND
 - Selective Medial Brachial Cutaneous Nerve (MBCN) and Intercostobrachial Nerve (ICBN) blocks
- Efficacy of Ultrasound-Guided (USG) VS Conventional Block of the MBCN and ICBN

Materials and Methods :

- Single-blinded randomized clinical trial in patients undergoing upper limb surgery, under locoregional anesthesia (1% Mepivacaine), at CHRU Nancy, between November 2016 and February 2017.
- Ultrasound-Guided Axillary brachial plexus block for all patients AND :
 - either USG (n=42) or conventional (n=42) block of the MBCN and the ICBN
- Light touch test at 5, 10, 15, 20 min in 4 areas : upper and lower half of the antero- and posteromedial surfaces of the upper arm
- Primary outcome : percentage of patients with no sensation in all these 4 areas at 20 min

Results :

- 42 patients per group
- In all patients : regional anesthesia alone

- Time to complete anesthesia : ≤ 10 min (in more than 90% of cases) in the 2 groups
- Mean total volumes of Mepivacaine for MBCN and ICBN blocks : 5 mL (USG group) VS 4,83 mL (Conventional group), $p = 0,334$
- Ultrasound images of good quality in 20 of 42 patients (47,6%) in the USG group
- Tolerance of tourniquet : comfortable in 41 patients in the USG group (97,6%) VS 16 patients (38,1%) in the Conventional group, $p < 0,01$

Conclusion :

- Ultrasound guidance improved the efficacy of the MBCN and ICBN blocks.