Risk Stratification of Thyroid Cancer using Thyroid Nodule Ultrasound Classification Systems: A Review Paper

Rakesh Mistry¹, Thushanth Sooriyaamoorthy², Waqas Din³, Nirmal Kumar⁴

¹ Airedale Hospital NHS Foundation Trust
² University of Birmingham Medical School
³ Hull and East Yorkshire NHS Trust
⁴ Warrington, Wigan and Leigh NHS Foundation Trust

Background:

Thyroid nodules are present in approximately 8% of the adult population on palpation, 41% on ultrasound scan and 50% on autopsy. Approximately 10% of these nodules are malignant³.

Investigation of suspicious nodules requires risk stratification via ultrasound scanning with a view to subsequent potential Fine Needle Aspiration Biopsy (FNAB). Currently there remains multiple systems utilised worldwide for risk stratification of thyroid nodule features on ultrasound scanning.

Objectives:

- (1) Review current evidence for ultrasound classification systems of thyroid nodules and their correlation to subsequent FNAB
- (2) Provide suggestions for future research.

Methods:

PUBMED and SCOPUS databases were searched in September 2017

Terms: ‘ultrasound classification’ AND ‘thyroid nodules’ AND ‘fine needle aspiration’ AND ‘malignancy’

Inclusion Criteria: research published in the last 5 years

Exclusion Criteria: If abstracts were irrelevant to the aims they were excluded.

This provided a total of 66 papers.

- 46 papers were then excluded as two authors assessed the titles of these papers for relevance to the aims
- 6 full papers were not accessible.

This resulted in a total of 14 papers for review

References

1. Singaporewalla RM, Hwee J, Lang TU and Desai V. Clinico-pathological Correlation of Thyroid Nodule Ultrasound and Cytology Using the TIRADS and Bethesda Classifications. World J Surg. 2017 Jul; 41(7): 1807-1811.
2. Middleton WD, Teefey SA, Reading CC, Langer JE, Beland MD, Szabunio MM, Desser TS. Multiinstitutional Analysis of Thyroid Nodule Risk Stratification Using the American College of Radiology Thyroid Imaging Reporting and Data System. AJR Am J Roentgenol. 2017 Jun; 208(6): 1331-1341
3. Rahal AJ, Falsarella PM, Rocha RD, Lima JP, Iani MJ, Vieira FA, et al. Correlation of Thyroid Imaging and Reporting Data System (TI-RADS) and fine needle aspiration: experience in 1,000 nodules. Einstein (Sao Paulo). 2016 Apr-Jun; 14(2): 119-23
4. Yoon JH, Lee HS, Kim EK, Moon HJ, Kwak JY. Malignancy Risk Stratification of Thyroid Nodules: Comparison between the Thyroid Imaging Reporting and Data System and the 2014 American Thyroid Association Management Guidelines. Radiology. 2016 Mar; 278(3): 917-24.
5. Chandramohan A, Khurana A, Pushpa BT, Manipadam MT, Naik D, Thomas N, et al. Is TIRADS a practical and accurate system for use in daily clinical practice? Indian J Radiol Imaging. 2016 Jan-Mar; 26(1): 145-52.
6. Rosário PW, Calsolari MR. Thyroid nodules with highly suspicious ultrasonographic features, but with benign cytology on two occasions: is malignancy still possible? Arch Endocrinol Metab. 2016 Aug; 60(4): 402-4.
7. Zhang J, Liu BJ, Xu HX, Xu JM, Zhang YF, Liu C, et al. Prospective validation of an ultrasound-based thyroid imaging reporting and data system (TI-RADS) on 3980 thyroid nodules. Int J Clin Exp Med. 2015 Apr 15; 8(4): 5911-7.

Results:

Three classification systems were identified in the literature:

- (A) Thyroid Imaging Reporting and Data System (TIRADS)
- (B) American Thyroid Association (ATA) Guidelines
- (C) Bethesda Classification System

Eight of the fourteen papers investigated TIRADS:

TIRADS 1	Normal
TIRADS 2	Benign
TIRADS 3	Probably benign
TIRADS 4	Suspicious
TIRADS 5	Probably malignant

Summary of Statistical Measurements from Literature for TIRADS:

Statistical Measurement	Range
Sensitivity	70.6 - 97.4%
Specificity	29.3 – 90.4%
Positive Predictive Value (PPV)	23.3 – 49%
Negative Predictive Value (NPV)	88 – 99%

Only Yoon et al. directly compared TIRADS to the ATA systems ($p < 0.001$ for all **PURPLE**):

Statistical Measurement	TIRADS	ATA
Sensitivity	97.4%	95.3%
Specificity	29.3%	37.4%
PPV	23.3%	98.1%
NPV	98.1%	97.3%
Accuracy	41.6%	47.9%

Yoon et al. reported a small proportion of nodules that could not be classified using the ATA criteria. These reported a similar malignancy rate to the intermediate risk group.

Conclusions:

- Current literature focuses heavily on TIRADS.
- The high NPV suggests TIRADS is more useful in ruling out malignancy whilst the low PPV may result in an increased number of unnecessary FNAB.
- Some nodules may not fall into ATA criteria and these may carry a relatively high malignancy risk
- Further research is required to directly compare different classification systems, particularly the British Thyroid Association guidelines here in the UK

8. Puno-Ramos MPG, Villa ML, Kasala RG, Arzadon J, Alcazaren EAS. Ultrasound Features of Thyroid Nodules Predictive of Thyroid Malignancy as Determined by Fine Needle Aspiration Biopsy. Philippine Journal of Internal Medicine. 2015 April-June; 53(2)
9. Adamczewski Z, Lewiński A. Proposed algorithm for management of patients with thyroid nodules/focal lesions, based on ultrasound (US) and fine-needle aspiration biopsy (FNAB): our own experience. Thyroid Res. 2013 Apr 20; 6(6).
10. Acharya UR, Faust O, Sree SV, Molinari F, Suri JS. ThyroScreen system: high resolution ultrasound thyroid image characterization into benign and malignant classes using novel combination of texture and discrete wavelet transform. Comput Methods Programs Biomed. 2012 Aug; 107(2): 233-41.
11. Gharib H, Papini E, Garber JR, Duick DS, Harrell RM, Hegedüs L. AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS, AMERICAN COLLEGE OF ENDOCRINOLOGY, AND ASSOCIAZIONE MEDICI ENDOCRINOLOGI MEDICAL GUIDELINES FOR CLINICAL PRACTICE FOR THE DIAGNOSIS AND MANAGEMENT OF THYROID NODULES--2016 UPDATE. Endocr Pract. 2016 May; 22(5): 622-39.
12. Yoon JH, Cho A, Lee HS, Kim EK, Moon HJ, Kwak JY. Thyroid incidentalomas detected on 18F-fluorodeoxyglucose-positron emission tomography/computed tomography: Thyroid Imaging Reporting and Data System (TIRADS) in the diagnosis and management of patients. Surgery. 2015 Nov; 158(5): 1314-22.
13. Papini E, Pacella CM, Hegedus L. Diagnosis of endocrine disease: thyroid ultrasound (US) and US-assisted procedures: from the shadows into an array of applications. Eur J Endocrinol. 2014 Mar 14; 170(4)
14. Heller MT, Gilbert C, Ohori NP, Tublin ME. Correlation of ultrasound findings with the Bethesda cytopathology classification for thyroid nodule fine-needle aspiration: a primer for radiologists. AJR Am J Roentgenol. 2013 Sep; 201(3)