[2+2] Cycloaddition of *p*-Toluenesulfonyl Isocyanate to Hydrocarbon Alkenes and Monofluoroalkenes

Francesca Flores, Dr. Dale Shellhamer, Ph.D

Department of Chemistry, Point Loma Nazarene University, 3900 Lomaland Drive, San Diego, Ca. 92106-2898

Abstract

We investigated the addition of *p*-toluenesulfonyl isocyanate (TsNCO) to electron-rich hydrocarbon alkenes and monofluoroalkenes to determine their ability to produce monofluoro-tosyl- β -lactam rings, and to determine if the reaction occurs via a Single Electron Transfer pathway.

β -lactam rings are a common structural motif found in antibacterial agents and potential cholesterol-lowering drugs.

Introduction

The β -lactam ring is found in medicinal chemistry studies. There are only a few successful [2+2] cycloaddition reactions reported with *p*-toluenesulfonyl isocyanate (**1**) to give the tosyl- β -lactam products since the four-membered ring opens at the high temperatures needed for reaction of alkenes with this less reactive electrophile. In an earlier study we found that neat reaction of isocyanates with alkenes proceed at lower temperatures and give better β -lactam product yields than reactions carried out in solvent.¹ In this study we treat **1** with alkenes in the absence of solvent to give tosyl- β -lactams (Scheme 1).

Recently also we reported on the use of line-broadening of an internal standard in the NMR to detect 1,4-diradical intermediates from the reaction of chlorosulfonyl isocyanate with alkenes and monofluoroalkenes in solution.² In this study we report on the reaction of 3,4-dihydro-2H-pyran (**5**) with **1**. The vinyl ether **5** was utilized because it is reactive enough to give a tosyl- β -lactam product in solution with the sluggish electrophile **1**, and allow for a search of the 1,4-diradical intermediate described in (Scheme 1).

Results & Conclusion

Alkenes were treated neat with Ts-N=C=O (**1**) at 50 °C. Products were isolated by column chromatography with ethyl acetate/hexanes as the eluant. Isolated product yields were **2** (59 %); **3** (58 %) and **4** (67 %) indicating successful reactions under neat conditions.

Line-broadening was observed when **1** was treated with the vinyl ether **5** in chloroform-d at room temperature (Compare Figure 1 and 2). Figure 3 shows that the NMR resolution was not lost during the reaction since the peak width at half-height returned to its original value when the reaction was complete. Line-broadening did not disappear at lower temperatures (Figure 4) as found for reaction of chlorosulfonyl isocyanate with electron-rich alkenes indicating a change from the Triplet to Singlet species.² It appears that the intermediate described in Scheme 1 remains a Triplet even at -10 °C.

Experimental

Line-broadening experiment. To a dry NMR tube was added 0.5 mL CDCl₃ with TMS as internal standard followed by addition of 0.12 mmol **5**. The initial spectrum was recorded and the TMS peak was expanded from -0.050 to +0.050 ppm to obtain the peak width at half-height. Ts-N=C=O (0.13 mmol) was added and the spectrum recorded without further shimming. TMS was analyzed as above. When the reaction was complete the TMS peak was measured again without shimming to confirm that resolution was not lost during the reaction.

References & Acknowledgements

1. Shellhamer, D. F.; Alexander, K. L.; Bunting, S. A.; Elwin, S. L.; Licata, C. J.; Milligan, J. C.; Robinson, R. D.; Shipowick, D. E.; Smith, L. B. and Perry, M. C. "Improved synthetic utility of a sluggish electrophile: Reaction of chlorosulfonyl isocyanate with unreactive and reactive alkenes", *Synthesis*, 2015, 47, 1994-1950.
2. Shellhamer, D. F.; Beavis, Z. J.; Brady, D. L.; Bucardo, M. S.; Elwin, S. L.; Fiorella, N.; Gomez, L.; Van Horne, S. and Perry, M. C. "NMR line-broadening and transverse relaxation time measurements support a di-radical intermediate for the reaction of chlorosulfonyl isocyanate with electron-rich alkenes" *Results in Chemistry* 2, (2020), 100015. <https://doi.org/10.1016/j.rechem.2019.100015>

Research Associates of PLNU, our alumni support group, provided support for this work.