P-0831– Association between Socioeconomic Status and Risk of Incident Diabetes in Adults Residing in Urban Delhi

Kanika Agarwal¹, Roopa Shivshankar², Mohammed K Ali³, KM Venkat Narayan³, Dorairaj Prabhakaran², Ravinder Chadha¹, Nikhil Tandon⁴

¹Department of Food and Nutrition, Lady Irwin College, University of Delhi, India
²Centre for Chronic Disease Control, Public Health Foundation of India, New Delhi, India
³Rollins School of Public Health, Emory University, Atlanta, United States of America
⁴Department of Endocrinology and Metabolism, All India Institute of Medical Sciences, New Delhi, India

Background

In developed countries, a lower socioeconomic status (SES) is associated with an increased risk of diabetes, while, in developing countries such as India, affluent people are found to be at a higher risk. With increasing urbanization, rural-to-urban migration, and change in lifestyle (*e.g.*, high calorie diet, physical inactivity, etc.), the risk of diabetes is also increasing among people in lower SES.

Aim

To assess the association between socioeconomic status and risk of incident diabetes in adults.

Methods

Study Design and Locale- Matched case control study, within the Cardio-metabolic Risk Reduction in South Asia (CARRS) cohort, a longitudinal study among non-pregnant adults ≥ 20 years, representative of urban Delhi. Data from the study were available from baseline evaluation and an annual follow up for the first two years, with clinical and biochemical evaluation performed at baseline and after 2 years; and only a clinical evaluation at year 1 of follow up.

Sample Size- 100 adults newly diagnosed with diabetes (fasting blood glucose ≥ 126 mg/dl or HbA_{1c} $\geq 6.5\%$) and 200 age and gender matched controls (diabetes free individuals at baseline and till the second annual visit)

Data Collected-

- *Socio economic status-* was assessed by educational qualifications, occupation, and income of the participants
- *Anthropometric measurements-* Height (stadiometer), Weight (weighing scale), and Body Mass Index (BMI)
- *Questionnaire-* Family history of diabetes, tobacco and alcohol consumption
- *International Physical Activity Questionnaire (IPAQ- short form)*

Statistical Analysis-

- Analyzed data using Stata version 11 and a 5% level of significance was considered
- Applied univariate conditional logistic regression analysis
- Applied conditional logistic regression with and without adjustment for covariates

Results

The difference in distribution of education, occupation, and income classes across cases and controls were found to be statistically insignificant

Table 1: Association between SES Status and Risk of Type 2 Diabetes

Variables	Model 1 [#]		Model 2 [#]	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Educational Qualification				
Primary school and below	0.952 (0.445, 2.035)	0.8981	1.400 (0.402, 4.871)	0.5969
High or secondary school	1.246 (0.700, 2.217)	0.4555	1.561 (0.630, 3.870)	0.3365
Graduate or higher	1.000		1.000	
Family Income (Rs/month)				
<10,000	0.682 (0.316, 1.470)	0.3290	0.856 (0.242, 3.029)	0.8097
10,000-30,000	1.199 (0.594, 2.421)	0.6127	1.532 (0.510, 4.606)	0.4472
30,000-50,000	1.195 (0.485, 2.943)	0.6989	1.854 (0.528, 6.510)	0.3351
$\geq 50,000$	1.000		1.000	
Occupation				
Not working	1.679 (0.718, 3.922)	0.2316	1.407 (0.396, 4.996)	0.5977
Semi or unskilled worker	1.049 (0.482, 2.285)	0.9043	2.444 (0.676, 8.835)	0.1729
Skilled worker	1.400 (0.600, 3.264)	0.4360	2.217 (0.659, 4.58)	0.1982
White collar	1.000		1.000	

[#]Model 1- individual univariate crude model for each variable, Model 2: Multivariate model, adjusted for body mass index (continuous), family history of diabetes (yes vs no), physical activity score (continuous), duration of tobacco use (continuous) and alcohol consumption (yes vs no)

Discussion

Our results show that the lower educational attainment / income, and/or non-white collar occupation is associated with risk of diabetes, similar to observed in the more affluent socio economic group. Until recently, the affluent in the country were thought to be at a higher risk of diabetes. However, we find that the risk of diabetes is becoming independent of the socioeconomic status. Therefore, the healthcare policies to prevent diabetes should evenly focus across all strata of the society.