Targeted drug therapy to provide effective and localised anticoagulation without impairing haemostasis

Xiaowei Wang^{1,2}, Donny Hanjaya-Putra³, Carolyn Haller³, Amy Kate Searle¹, Karlheinz Peter^{1,2}, Elliot L. Chaikof³

¹Atherothrombosis and Vascular Biology, Baker Heart & Diabetes Institute, Melbourne, Victoria, Australia; ²Department of Medicine, Monash University, Victoria, Australia; ³Beth Israel Deaconess Medical Center (BIDMC), Harvard Medical School, Boston, Massachusetts

INTRODUCTION

- Current dual combination therapy using anti-platelet; anti-thrombotic; and/or anti-coagulation agents offers the promise of treatment and improved thrombotic protection
- Substantial risk of major bleeding hampers widespread use

METHODS

- Single chain antibody
- Targeting blocks activated GPIIb/IIIa complex on activated platelets (scFv^{Targ})
- Non-binding mutated control (scFv^{Non-targ})
- Tick anticoagulant peptide (TAP)
- Potent direct inhibitor of FXa
- scFv^{Targ} + TAP = Targ-TAP
- scFv^{Non-targ} + TAP = Non-targ-TAP

RESULTS

Targ-TAP binds to activated platelets and retains anti-FXa activity after fusion

Targ-TAP prevents thrombosis *in vitro*

Targ-TAP colocalises with CD42b

Targ-TAP increases time to occlusion without bleeding *in vivo*

Subcutaneous prophylaxis approach (arterial)

Subcutaneous prophylaxis approach (venous)

Targ-TAP binds after sealing layer of platelets

CONCLUSION

Activated platelet targeted TAP

- Prevents thrombosis *in vitro* and *in vivo*
- Overcomes bleeding complications
- Enables prophylactic strategy to be employed

xiaowei.wang@baker.edu.au