Background

The experience of a forced migration is a risk factor for the maintenance and restoration of mental health, leading to an increased prevalence of post-traumatic stress, depression and anxiety disorders (Turrini et al., 2017; Boden et al., 2012). In the course of the so-called *refugee crisis*, many German communities initiated programs that focused on musical activities with refugees assumed to be beneficial to the refugee's integration. However, the effectiveness of such music-centered programs has not yet been investigated.

Objective: The present study aims at investigating effects of musical activity and music-making on psychological adaptation in adult male refugees.

Methods

Sample: N = 87 ($M_{age} = 23.4$ yrs, $SD_{age} = 4.17$, Range_{age} = 18 – 34 yrs) adult male refugees; musically untrained.

Music intervention: Six months of weekly music sessions in small groups (approx. 2 hrs), led by experienced musicians.

Measures:

General Health Questionnaire (GHQ): Self-assessment of mental health ($\alpha = .85 - .91$).

Connor-Davidson-Resilience-Scale (CD-RISC): Psychological resilience ($\alpha = .87 - .91$).

PTSD- Checklist for DMS-V (PCL-5): PTSD-symptoms ($\alpha = .85 - .91$).

Perceived Stress Scale (PSS): Perceived stress levels ($\alpha = .83 - .85$).

Frankfurt Acculturation Scale (FRACC): Orientation Towards Host Culture (HC; $\alpha = .48 - .67$), Orientation Towards Culture of Origin (OC; $\alpha = .55 - .67$).

Emotion Regulation Questionnaire (ERQ): Reappraisal of emotions (ERQ-ReApp, $\alpha = .72 - .84$).

Study Design

Handling of critical drop-out rates (~40%): Multiple imputation of missing values (Demir et al., 2016; Sullivan et al., 2018)

Results

Analysis of longitudinal data (completed data set) using linear mixed effects models (*lmem*)

Changes over time in health-related outcomes

Changes over time in acculturation outcomes

Changes over time in emotion regulation and resilience

Discussion

Data analysis suggests that music activities in groups with refugees may lead to a substantial decrease in symptoms of post-traumatic stress, stress experience and an increase in the orientation towards host culture. Effects were observed after a time interval of 6 months between pre- and post-measurement.

Jasmin Chantah, chantah@psych.uni-frankfurt.de Prof. Dr. Stephan Bongard, bongard@psych.uni-frankfurt.de