Chemotherapy-induced Neutropenic Septic Shock in Gynecologic Cancer : Clinical course and characteristics

Ji-Geun Yoo, Yoon-Jin Choi, Jong-Sup Park, Keun-Ho Lee, Soo-Young Hur

Department of Obstetrics and Gynecology, Seoul St. Mary's Hospital,
The Catholic University of Korea

Introduction

For advanced gynecologic cancer patients, Chemotherapy has provided survival benefits to cancer patients, but at the same time has a risk of bone marrow suppression and possibility of combined infection, and occasionally cause lethal septic shock, which is associated with a mortality rate of 40%. Still, there is only a few reports about chemotherapy-related neutropenic septic shock in gynecologic malignancies. The purpose of this study is to assess the risk and determine clinical characteristics and course of septic shock in gynecologic cancer patients.

Methods

In this single-center retrospective study we reviewed medical records of gynecologic cancer patients who had treated with chemotherapeutic agents between Jan. 2009 and Dec. 2017. Neutropenic septic shock is defined as neutropenia <1g/L with septic shock that requires inotropics to maintain normal blood pressure. We divided the events into two groups according to survival. Statistical analysis was performed for the clinical characteristics by Mann-Whitney U-test.

Results

Total 1363 patients received at least 1 cycle of chemotherapy for gynecologic malignancies. Among them, 22(1.6%) patients experienced 25 events of neutropenic septic shock. Six events of death have occurred, and the mortality ratio was 24%.

Table 1. Clinical characteristics of the events.

Characteristics	Total	Events of survive	Events of death	P value
Events, n	25	19 (76.0%)	6 (24.0%)	
Median age, yrs	59 [29-78]	57 [29-66]	62 [55-78]	.085
Mean BMI, kg/m ²	22.9 [14.7-28.4]	22.9 [14.7-28.4]	22.7 [18.2-27.5]	.874
Type of cancer				
Ovarian	13 (52.0%)	9 (47.4%)	4 (66.7%)	
Endometrial	6 (24.0%)	4 (21.1%)	2 (33.3%)	
Peritoneal	3 (12.0%)	3 (15.8%)		
Cervical	2 (8.0%)	2 (10.5%)		
Uterine leiomyosarcoma	1 (4.0%)	1 (5.3%)		
FIGO Stage, n(%)				
I	1 (4.0%)	1 (5.3%)	0	
II	4 (16.0%)	3 (15.8%)	1 (16.7%)	
III	11 (44.0%)	7 (36.8%)	4 (66.7%)	
IV	9 (36.0%)	8 (42.1%)	1 (16.7%)	
Recurrence status				
Primary treatment, n(%)	8 (32.0%)	6 (31.6%)	2 (33.3%)	
Recurrent disease, n(%)	17 (68.0%)	13 (68.4%)	4 (66.7%)	
Duration from the initial chemotherapy(mo.)	23 [1-131]	18 [1-85]	28 [1-131]	.799
Days from the latest chemotherapy(d.)	8 [3-14]	8 [3-14]	8 [4-14]	.923
Peak body temperature*(°C)	38.8 [35.6-40.4]	39 [37.3-40.4]	38.3 [35.6-40.4]	.251
Mean absolute neutrophil count*, mg/L	150 [0-780]	122.6 [0-550]	236.7 [0-780]	.305
Mean CRP at diagnosis, mg/dL	17.8 [1.3-30.4]	17.4 [1.28-29.1]	18.9 [9.49-30.4]	.703
Median duration of neutropenia, days	3 [1-7]	4 [1-7]	3 [1-6]	
Mean duration of inotropics, Hr	50.6 [6-191.9]	49.3 [6-140.8]	54.8 [11-191.9]	.899

*The lowest or highest value during event

There was no significant difference between events of death group and survive group according to age, BMI, FIGO stage, mean CRP at the time of diagnosis of septic shock, the lowest neutrophil count during the event, and peak body temperature.

Six events occurred during the primary treatment course. We experienced two events of septic shock who received chemotherapeutic agents only once. According to the chemotherapeutic regimen, carboplatin-paclitaxel combination chemotherapy accounted for the largest proportion(24%).

Figure 1. Chemotherapy regimen related to septic shock.

Two-thirds of events had identified bacteria on blood culture. Most common pathogen isolated was E. coli(7/16, 43.7%).

In the event of survivor group, mean recovery time was 2 days of initial treatment. Meanwhile, for events of death group, the average time from initial treatment of septic shock to death was 54.8 hours.

Table 2. Identified pathogen on blood culture

	Total(n=25)	Events of survive(n=19)	Events of death(n=6)
E. Coli	7 (28.0%)	5 (26.3%)	2 (33.3%)
ESBL(-)	5 (20.0%)	3 (15.8%)	2 (33.3%)
ESBL(+)	2 (8.0%)	2 (10.5%)	0
Enterobacter Cloacae	2 (8.0%)	2 (10.5%)	0
Klebsiella Pneumoniae	3 (12.0%)	2 (10.5%)	1 (16.7%)
Pseudomonas Aeruginosa	1 (4.0%)	0	1 (16.7%)
Enterococcus Faecalis	1 (4.0%)	0	1 (16.7%)
Others	2 (8.0%)	2 (10.5%)	0
Not identified	9 (36.0%)	8 (42.1%)	1 (16.7%)

Conclusion

Through this study, we confirmed the incidence and mortality, and clinical course of neutropenic septic shock in gynecologic malignancy. We hope that this study will provide clinicians with information about chemotherapy related septic shock and help patients recover from this lethal disease.