Impact on outcome of personality disorders in recurrent major depressive disorder

J. Nakhli, Y. El kissi, S. Ben Nasr,

Farhat Hached University Hospital, Sousse, Tunisia

Background:

In literature, comorbidity between depressive disorder and personality disorder is common and varies from 18 to 51% [1]. It’s often responsible of impaired outcome and makes difficulties in treatment [2,3].

Aims:

The aim of this study was to determine the impact of personality disorders on the clinical characteristics and evolution of recurrent major depressive disorders.

Methods:

This is a retrospective descriptive study. It concern 211 female inpatients with recurrent depressive disorders according to DSM-IV during the period from 1999 to 2016. All these patients were admitted in psychiatric unit in Farhat Hached General Hospital (Sousse) because of depressive symptoms and/or suicidal ideation or attempts.

Exclusion criteria were: a diagnosis of bipolar disorder, other psychotic disorders and mental retardation.

All these female inpatients were followed up for a period of 24 months.

To evaluate impact of personality disorders on outcome of these female inpatients with recurrent major depressive disorder, we have compared two groups:

- First group (G1): inpatients with recurrent depression and having comorbid personality disorder (N = 120).
- Second group (G2): inpatients with recurrent depression without personality disorder (N = 91).

Statistical comparisons were performed between the two groups and based on demographic, clinical and outcome features available in patients’ medical charts.

Comparisons were carried out with χ^2 and t-test statistics. Reported differences were significant at 0.05 or less.

Results:

The mean age of our sample was 43.3 ± 13.1 years. They were married in 61% of cases and third of them had regular professional activity.

44% of our inpatient with recurrent depression had history of organic pathology and 34% of them had other comorbid psychiatric illness such us generalized anxiety disorder.

The mean age of onset of depressive disorder was 31.5 ± 11.5 years.

The mean number of admission in psychiatric unit was 2.1 ± 1.1 .

The prevalence of personality disorder in patients with recurrent major depressive disorders was 56.9%. The most frequent was respectively histrionic (40.8%), dependant (16.6%) and borderline (15.8%).

In the group of patients with recurrent depression and having personality disorder, the beginning of the first depressive episode and age of first medical visit and admission in psychiatry were earlier ($p < 10^{-3}$). Similarly, we found they had more suicide attempts in their history ($p < 10^{-3}$).

After a period of follow up, we found that they have more residual symptoms ($p = 0.01$), a larger number of relapses ($p = 0.042$) and subsequent suicidal recurrence ($p = 0.001$) (**Table I**).

Discussion:

Clinical factors

Depressed inpatients with personality disorders, in our study, had lower age at the onset of depressive illness than patients without PD. This result is noticed in many previous studies [2,3,4]. According to Ramklint and al., early onset of depression is a major predictor for personality comorbidity [4].

Personality disorders were associated with more suicide attempts in our depressed patients, as it was regularly noticed by authors [5,6]. Cluster B personality types, particularly borderline and antisocial ones, were the most prevalent in depressive patients with suicide attempts [5,6,7]. Many data assumed that specific traits of these personality types, such as impulsivity, hostility and aggressiveness may facilitate suicidal behavior [5,6,7]. Hansen and al. notified that PD favor the repetition of suicidal behavior [8].

Many studies had found that PD were associated with many axis I comorbid disorders like alcohol and substance abuse [2,3]. Joyce and al. reported that depressed patients with PD had significantly more alcohol and cannabis abuse or dependence and social phobia than those without PD [9].

Outcome factors

Feske and al. noticed that patients with MDD and PD had more partial remission and more relapses [10]. In the same way, Shea and al. had reported a more residual depressive symptoms in depressed patients with PD and impaired outcome in social functioning [11].

About recurrence, Alnaes and al. admitted that PD were responsible of recurrence in depression [12]. This result was contested by Joyce who didn’t find this association [9].

Conclusion:

Almost half of our inpatients followed for recurrent depressive disorders have associated personality disorder. The most frequent was respectively histrionic, borderline and dependant personality.

This comorbidity is associated with earlier age of first depressive episode, more suicide attempts, residual symptoms and relapses. Standardized assessment of personality disorder should be used in clinical practice to ensure adequate care. Other multicenter studies recruiting more patients with major depressive disorders are needed.

Table I. Clinical factors associated in personality disorder

	Group I	Group II	p
Age at onset of depressive illness (years)	28.1	34.8	<10 ⁻³
first admission in psychiatry (years)	29.9	37.6	<10 ⁻³
Past suicide attempts (%)	43	21	<10 ⁻³
Persistant depressive symptoms (%)	44	34	0.01
Relapses (%)	33	18	0.042
Subsequent suicidal recurrence (%)	9	3	0.001

No conflict of interest

REFERENCES

1- Newton-Howes G, Tyrer P, Johnson T. Personality disorder and the outcome of depression: meta-analysis of published studies. Br J Psy 2006; 188: 13-20.

2- Boyce P, Parker G, Barnet B and al. Personality as a vulnerability factor to depression. Br J Psy 1991; 159: 106-14.

3- Black DW, Bell S, Bulbert J and al. the importance of axis II in patients with major depression. A controlled study. J Aff Disor 1988; 14: 115-22.

4- Ramklint M, Ekselius L. Personality traits and personality disorders in early onset versus late onset major depression. J Aff Disor 2003; 75: 35-42.

5- Corbitt EM, Malone KM, Haas GL. Suicidal behavior in patients with major depression and comorbid personality disorders. J Aff Disord 1996; 39:61-72.

6- Gunderson JG, Phillips KA. A current view of the interface between borderline personality disorder and depression. Am J Psy 1991;148:967-75.

7- Oquendo M, Bongiovi-Garcia M, Galfalry H, Goldberg P, Grunebaum M. Prospective study of clinical predictors of suicidal acts after a major depressive episode in patients with major depressive disorder of bipolar disorder. Am J Psy 2004;1433-41.

8- Hansen PE, Wang AG, Stage KB, Kragh-Sorensen P. comorbid personality disorders predicts suicide after major depression: a 10-year follow-up. Acta Psychiat Scand 2003, 107(6): 436-40.

9- Joyce PR, Mulder RT, Luty SE and al. borderline personality disorder in major depression: symptomatology, temperament, character differential drug response and 6 months outcome. Compr psy 200"; 44(1): 35-44.

10- Feske U, Mulsant BH, Pilkonis PA. Clinical outcome of ECT in patients with major depression and comorbid personallity disorder. Am J Psy 2004; 161: 2073-80.

11- Shea MT, Pilkonis PA, Beckham E, Collins JF and al. Personality disorders and treatment outcome in the NIMH treatment of depression collaborative research program. Am J Psy 1990; 147: 711-18.

12- Alnaes R, Torgersen S. Personality and personality disorder predict development and relapse of major depression. Acta Psy Scan 1997; 95: 336-42.