Service Evaluation of the Use of the Young Person's Proforma

Aim: To investigate the adherence to BASHH guidelines relating to the care of young people accessing sexual health services, specifically the use of the Young Persons' Proforma.

¹Megan O'Connell, ²Nicola Lomax. ¹Cardiff University, Cardiff; ²Department of Sexual Health, Cardiff Royal Infirmary

Introduction

17% of women and 24% of men have sexual intercourse before the age of 16¹. Although consensual in some cases, the prevalence of non-volitional sex was higher in those with a sexual debut before age 16 years². National guidelines set by BASHH³ (British Association of Sexual Health and HIV) advocate the use of a Young Person's Proforma (YPP) in sexual health clinics to detect signs of, and concurrent risk factors for, child sexual exploitation (CSE), intending to safeguard this vulnerable group of attendees.

Methods

Discussion

- CRI achieved all BASHH targets, whereas the community clinics failed to do so
- There is a disparity in male attendance and subsequent proforma completion. Methods to effectively target young men need to be explored
- Incorporating SERAF scoring into the proforma effectively identifies those at risk of child sexual exploitation
- Outcomes of community referral were vague, indicating a disparity in communication between central and community clinics

Recommendations

- Placement of a proforma into notes before consultation
- Introducing a standardised time-frame for reassessment, along with a method of highlighting this within the notes
- Adopting a more opportunistic approach to STI testing in the community
- Introducing electronic case records, to aid communication and improve patient safety.
- Consider widening proforma usage to all patients under 18

Conclusion

Discrepancies in the use of the YPP between central and community clinics are evident, this may be partly due to the pressures of the service provision in the community. However, maintaining a consistent approach to the assessment of safeguarding between sites is of upmost importance, as is implementing other improvements that facilitate the creation of an equitable service.

References

(1) Mercer C, Tanton C, Prah P, Erens B, Sonnenberg P, Clifton S et al. Changes in Sexual Attitudes and Lifestyle in Britain Through the Life Course and Over Time: Findings from the National Surveys of Sexual Attitudes and Lifestyles (NATSAL). Lancet. 2013; 382: 1781-94.

(2) Macdonald V, Gibson L, Tanton C, Mercer C, Lewis R, Clifton S et al. Lifetime Prevalence, Associated Factors, and Circumstances of Non-Volitional Sex in Women and Men in Britain: Findings From the Third National Survey Of Sexual Attitudes and Lifestyles (NATSAL-3). Lancet. 2013; 382:1845-55.

(3) Rogstad K, Thomas A, Williams O, Forster G, Munday P, Robinson A. United Kingdom National Guideline on the Management of Sexually Transmitted Infections and Related Conditions in Children and Young People. British Association of Sexual Health and HIV; 2010. [Accessed 15/06/2015]. Available from: <http://www.bashh.org/documents/2674.pdf>