Prediction of survival with the changing pattern of vital sign in terminally ill cancer patients

Se-Il Go¹, Jung Hun Kang²

¹Department of Internal Medicine, Gyeongsang National University Changwon Hospital, Gyeongsang National University School of Medicine, Institute of Health Sciences, Changwon, Republic of Korea
²Department of Internal Medicine, Gyeongsang National University School of Medicine, Institute of Health Sciences, Jinju, Republic of Korea

Background

Prediction of survival is of great concern to terminally ill cancer patients and their family. Inaccurate prediction of survival results in unnecessary and potentially harmful therapies, prolongs hospitalization, and hinders patients from planning their remaining life. Treating physicians often rely on symptomatic change to predict survival of patients and tend to overestimate the remaining survival. Several prediction tools and scoring systems developed to overcome the inaccuracy of clinical estimation are rather complex to use easily by physicians. Therefore, objective and simple prediction model is needed to identify the patients who benefit from early palliative care.

Methods

Design and period: retrospective study (Oct. 2009 ~ Dec. 2013)

Eligibility

- Patients with advanced cancer who died at the regional cancer center of Gyeongsang National University Hospital during the study period
- Not treated with chemotherapy during the last 6 weeks of life
- At least one or more vital signs [systolic blood pressure (SBP), diastolic blood pressure (DBP), heart rate (HR), respiratory rate (RR), body temperature (BT)] obtained during the last 8 days

Study objectives and definitions

- Changes in vital signs over time during the last 8 days of life
- Differences of the average vital signs between day 0 – 3 (within 3 days) and day 3 – 8 (over 3 days) before death
- Impact of ‘vital sign risks’ on survival prediction
- ‘Vital sign risk’ was considered to be present if the daily average change of vital sign compared to the average vital sign of day 3 – 8 (over 3 days) before death meets the following criteria: SBP decrease ≥ 10 mmHg, DBP decrease ≥ 10 mmHg, HR increase ≥ 10 BPM, RR increase ≥ 5 RPM, and BT increase ≥ 0.5 °C

Results

Risk of death within 3 days according to the vital sign risk

Vital sign risk	within 3 days	over 3 days		Matched OR (95% CI)
		present	Absent	
SBP dec. ≥10 mmHg	Present	259	601	6.5 (5.2 - 8.1)
	Absent	93	278	
DBP dec. ≥10 mmHg	Present	87	587	11.1 (8.4 - 15.0)
	Absent	53	504	
HR inc. ≥10 BPM	Present	261	517	5.9 (4.7 - 7.5)
	Absent	87	366	
RR inc. ≥5 BPM	Present	38	430	14.8 (10.2 - 22.4)
	Absent	29	734	
BT inc. ≥0.5 °C	Present	64	309	3.4 (2.7 - 4.3)
	Absent	91	767	
2 of 5 vital signs	Present	265	748	15.6 (11.6 - 21.3)
	Absent	48	170	
3 of 5 vital signs	Present	72	653	17.6 (12.7 - 25.3)
	Absent	37	469	
4 of 5 vital signs	Present	12	362	25.9 (15.2 - 47.8)
	Absent	14	843	
5 of 5 vital signs	Present	3	178	35.6 (15.0 - 111.0)
	Absent	5	1045	

Sensitivity, specificity, and positive predictive value for the probability of death within 3 days according to the vital sign risk

	Sensitivity (95% CI)	Specificity (95% CI)	Positive predictive value (95% CI)
SBP dec. ≥10 mmHg	0.699 (0.672-0.724)	0.714 (0.688-0.739)	0.902 (0.882-0.920)
DBP dec. ≥10 mmHg	0.548 (0.520-0.575)	0.886 (0.867-0.903)	0.927 (0.906-0.944)
HR inc. ≥10 BPM	0.632 (0.605-0.659)	0.717 (0.692-0.742)	0.899 (0.878-0.918)
RR inc. ≥5 BPM	0.380 (0.354-0.408)	0.946 (0.932-0.957)	0.942 (0.918-0.959)
BT inc. ≥0.5 °C	0.303 (0.278-0.329)	0.874 (0.854-0.892)	0.804 (0.765-0.838)
2 of 5 vital signs	0.823 (0.801-0.843)	0.746 (0.721-0.769)	0.955 (0.941-0.966)
3 of 5 vital signs	0.589 (0.561-0.616)	0.911 (0.894-0.926)	0.951 (0.934-0.965)
4 of 5 vital signs	0.304 (0.279-0.330)	0.979 (0.969-0.986)	0.964 (0.940-0.978)
5 of 5 vital signs	0.147 (0.128-0.168)	0.994 (0.987-0.997)	0.973 (0.939-0.989)

A total of 1,357 patients were included. Mean SBP and DBP abruptly decreased during the last day of life. Mean RR apparently increased since 2 days before death. Mean HR consistently increased and there was small difference of mean BT during the last 8 days of life. The matched odds ratios for death within 3 days were 6.5, 11.1, 5.9, 14.8, and 3.4 in SBP, DBP, HR, RR, and BT, respectively, compared with their counterparts. The positive predictive values for death within 3 days were 90.2%, 92.7%, 89.9%, 94.2%, and 80.4% in SBP, DBP, HR, RR, and BT, respectively. The combined use of vital sign risks increased the positive predictive value up to 97.3%.

Conclusion

The change patterns of vital signs are insensitive to predict the impending death within 3 days, but may be useful to differentiate it from temporary deterioration due to the high positive predictive value in terminally ill cancer patients. A prospective study is warranted to validate these results.