Conidial surface proteome analysis reveals the highly abundant CcpA protein important for virulence of *Aspergillus fumigatus*

V. Voltersen^{1,2}, M. G. Blango¹, S. Herrmann³, F. Schmidt¹, T. Heinekamp^{1,2}, M. Strassburger⁴, T. Krüger¹, P. Bacher⁵, J. Lothar⁶, E. Weiss⁶, K. Hünig^{7,8}, H. Liu⁹, P. Hortschansky^{1,2}, A. Scheffold⁵, J. Löffler⁶, S. Krappmann¹⁰, S. Nietzsche¹¹, O. Kurzai^{7,8}, H. Einsele⁶, S. G. Filler⁹, U. Reichard³, O. Kniemeyer^{1,2}, A. A. Brakhage^{1,2}

¹Department of Molecular and Applied Microbiology, Leibniz Institute for Natural Product Research and Infection Biology (HKI), Jena, Germany; ²Department of Microbiology and Molecular Biology, Institute of Microbiology, Friedrich Schiller University Jena, Jena, Germany; ³Institute for Medical Microbiology, University Medical Center, Göttingen, Germany; ⁴Transfer Group Anti-infectives, HKI, Jena, Germany; ⁵Department of Cellular Immunology, Clinic for Rheumatology and Clinical Immunology, Charité - University Medicine, Berlin, Germany; ⁶Medical Clinic and Policlinic II, University Clinic Würzburg, Würzburg, Germany; ⁷Institute for Hygiene and Microbiology, University of Würzburg, Würzburg, Germany; ⁸Septomics Research Center, HKI, Jena, Germany; ⁹Division of Infectious Diseases, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, and David Geffen School of Medicine at University of California, Los Angeles, California, USA; ¹⁰Microbiology Institute - Clinical Microbiology, Immunology and Hygiene, University Hospital Erlangen and Friedrich Alexander University of Erlangen-Nürnberg, Erlangen, Germany; ¹¹Electron Microscopy Centre, Clinics of the Hospital, Jena, Germany

Introduction

Aspergillus fumigatus is a common airborne fungal pathogen of humans and a significant source of mortality in immunocompromised individuals. Conidia, the asexually produced spores, are the first form of *A. fumigatus* to come in contact with lung epithelial and immune cells. Thus, the composition of the conidial surface has the potential to directly influence the outcome of an *A. fumigatus* infection. Here, we aimed to identify additional conidial surface proteins which may play a role during host pathogen interaction by either hydrogen fluoride-pyridine (HF/pyridine) extraction of conidial proteins or "trypsin-shaving" of surface-exposed proteins and subsequent LC-MS/MS analysis.

Experimental procedures

Conidial surface proteome

Results

The conidial cell wall protein CcpA is a highly abundant protein on the spore surface

```
> CcpA (A. fumigatus Af293)
1  MHYSQFVTVA SALALAPTAV VARQGAAAFV TVNSIDVCPK KVAQEIIINPG
51  PKVVTTPYTQ DQVKLGHLG DVSYYNFDIEP LTKDTFPYCK ALKVFDEGEC
101 LGFPTLWIPL ESPLDKCIP EHYFSDEVKS ISFQLDCRED APVKKEPYGP
151 KEGAEQSAPQ AEHSTKQDAQ QGSHQGQEVQ NSPKQEARQG SRPAEAAAPKQ
201 EQEAEQASEA APEKKASNPA DSLGLGELTK VLGFR
```

- CcpA is an uncharacterised conidial surface protein with signal peptide for secretion and 6 conserved cysteines
- It is present in *Aspergilli* and closely related species (*Penicillium*, *Neosartorya*)

The conidial cell wall protein CcpA is surface localised, conidia-specific, and shows developmental expression

A *ccpA* deletion-strain (Δ *ccpA*) does not show obvious growth phenotypes or changes of conidial surface properties

The surface proteome of Δ *ccpA* is altered as revealed by surface shaving of dormant conidia with trypsin

The deletion of *ccpA* leads to attenuated virulence in corticosteroid-treated mice and increased the oxidative burst in PMNs

Conclusions

CcpA is a surface-localised, structural spore protein of *A. fumigatus* that is important for virulence.