Assessing the validity of overnight sleep studies in paediatric patients in a district general hospital.

A Harrison, A Anwar, A Hobbs

Obstructive Sleep Apnoea is a disorder of breathing during sleep characterised by prolonged partial upper airway obstruction and/or intermittent complete obstruction that disrupts normal ventilation during sleep. This can be assessed with overnight oxygen saturations.

Children presenting with clinically significant sleep related breathing difficulties typically have their symptoms effectively managed by adenotonsillectomy

Methods

- The investigation of these children is variable and evidence regarding overnight oxygen saturation (ON) is contradictory

Objectives

- To assess the use of ON in our district general hospital (Royal Bolton Hospital) and its ability to predict postoperative complications.

Methods

- Retrospective analysis of 195 paediatric adenotonsillectomy procedures between January and December 2016.
- No formal guideline or guidance at RBH on who should have a planned overnight admission, whether it be HDU or ward, and what sort of monitoring was needed if admitted for OSA.
- Data collected included use of pre operative ON, severity score allocated, oxygen desaturation index (ODI) and post operative saturations especially any apnoeas.

Discussion

ODI score used alone does not adequately identify at risk patients and may underreport significant isolated low saturations (sats nadir) . Patient comorbidities as discussed in the 2009 consensus statement¹ and sats nadir seem more reliable indicator of post operative apnoeas.

Table 2
Children at risk from respiratory complications unsuitable for DGH adenotonsillectomy

Age <2 years
Weight <15kg
Failure to thrive (weight <5 th centile for age)
Obesity (BMI >2.5SDS or >99 th centile for age and gender)
Severe cerebral palsy
Hypotonia or neuromuscular disorders (moderately severely or severely affected)
Significant craniofacial anomalies
Mucopolysaccharidosis and syndromes associated with difficult airway
Significant comorbidity (e.g. congenital heart disease, chronic lung disease, ASA 3 or above)
ECG or echocardiographic abnormalities
Severe OSA (described by polysomnographic indices including Obstructive Index >10, Respiratory Disturbance Index >40, and Oxygen saturation nadir <80%)

Consensus statement ¹.

Results

- 195 cases [98 M , 97 F]
- Age 2 years to 15 years 6 months
- 53 Patients had sleep studies (27%)
 - Normal 41 (77%)
 - Mild 8 (15%)
 - Moderate 3 (6%)
 - Severe 1 (2%)
- 4 (2%) had an apnoeic episode post operative
 - 1 Normal sleep study / 3 none
 - 1<15KG; 1<9TH CENTILE; 1>99TH CENTILE
 - All had unplanned stay as none were identified as at risk pre operatively with current assessment using ODI alone.

Conclusion

ODI used alone does not adequately identify at risk patients and may underreport significant isolated low saturations. Patient comorbidities and sats nadir could be used to predict complications

1. Robb PJ, Bew S, Kubba H, Murphy N, Primhak R, Rollin AM, Tremlett M Tonsillectomy and adenoidectomy in children with sleep-related breathing disorders: consensus statement of a UK multidisciplinary working party. Ann R Coll Surg Engl. 2009 Jul;91(5):371-3