Effect of implementation of a specialised paediatric anaesthesia team and training programme on incidences of perioperative paediatric cardiac arrest: a before and after analysis.

A. Hohn, U. Trieschmann, J. Kaufmann, J. Hinkelbein, S. A. Padosch, B. W. Böttiger

Background and Goal of Study: This study analyses incidences of perioperative paediatric cardiac arrest (CA) before and after implementation of a specialised paediatric anaesthesia team and training programme at tertiary care university hospital in Germany.

Materials and Methods: Reports on perioperative paediatric (0-18 yrs) CA were identified from all critical incident reports 2008-2016. CAs were classified as anaesthesia-attributable or anaesthesia-unrelated. Incidences of CA were calculated from the anaesthesia database. A logistic regression analysis with risk factors (age, ASA physical status grade [ASA-PS], emergency) was used to evaluate the impact of implementation of a specialised paediatric anaesthesia team and training programme in 2014 on perioperative paediatric CA.

Results: Between 2008 and 2016, 25 perioperative paediatric CAs in 36,243 paediatric anaesthesias were identified. Of these, 12 were classified as anaesthesia-attributable CA. The incidence of anaesthesia-attributable CA was 4.3 (95%-CI 2.3-7.9) in the period 2008-2013 and decreased to 1.6 (95%-CI 0.3-5.7) in 2014-2016 (*Figure 1*). Likewise, the incidence of overall CA was lower after 2013

[4.6/10,000 (95%-CI 2.1-10.2) vs 8.1/10,000 (95%-CI 5.2-12.7)]. Children anaesthetised after 2013, had nearly a 70 percent lower probability of anaesthesia-attributable CA (OR 0.306; 95%-CI 0.067-1.397; $p=0.1263$) although proportions of children aged <1 year and classified ASA-PS 3-5 were greater in the period 2014-2016. For anaesthesia-attributable CA, young age was the most contributing risk factor whereas in overall paediatric CA, ASA-PS 3-5 played a more important role. During the study period, 91.7% (11/12) of anaesthesia-attributable CAs occurred in children < 3 years.

Conclusions: Young age, ASA-PS and cardiac procedures contribute as main risk factors to perioperative paediatric CA. In this study on the incidences of perioperative paediatric CA from a European tertiary care university hospital, implementation of a specialised paediatric anaesthesia team and training programme was associated with lower incidences of perioperative paediatric CA and a reduced probability for anaesthesia-attributable CA.

Figure 1 Annual incidences of perioperative CA and anaesthesia-attributable CA 2008-2016 (A/C), and incidences of CA before (2008-2013) and after (2014-2016) implementation of a specialised paediatric anaesthesia team (B/D). Proportions of children aged <1 yr and < 1 yr + ASA-PS 3-5 respectively (E).