QUALITY OF LIFE OF PATIENTS WITH TRACHEOSTOMY AFTER DISCHARGE

¹ÇIRACI YASAR Y, ¹NURAL N, ²UYAR Y, ²SALTURK Z

¹KARADENİZ TEKNİKAL ÜNİVERSİTESİ SAĞLIK BİLİMLERİ FAKÜLTESİ HASTALIK BİLİMİ ANA BİLİMİ

²OKMEYDANI TERCİME VE ARAŞTIRMA HASTAHANESİ KLİNİK BİLİMİ

Purpose: Patients with a permanent tracheostomy experience physical and emotional problems secondary to surgical procedures. The purpose of this study was to analyze QOL in patients with a permanent tracheostomy and the factors affecting their QOL.

Method: This is a cross-sectional and descriptive study. Forty-five patients underwent total laryngectomy at Okmeydanı Training and Research Hospital Ear Nose and Throat Clinic in 2015 and 36 of those discharged with permanent tracheostomy participated in this study. Sociodemographic data were collected along with data on QOL using EORTC QLQ C-30 [European Organization for Research and Treatment of Cancer (EORTC), Quality of Life (QOL) Group]. The nonparametric Kruskal Wallis and Mann Whitney U tests were used for the analysis of sociodemographic data and variables that could affect QOL. The significance level was accepted as 0.05.

Results: The mean common wellbeing and quality of life score was 47.6 ± 14.4 , which is below average. Regarding functional status, cognitive and emotional functions were affected the most. Regarding the symptom subscale, insomnia and fatigue were the factors influencing decreased quality of life the most. Moreover, age and social environment were major contributors of quality of life in the symptom subscale. With an increase of age and a decline in social environment, quality of life declines. Social environment also affects patients' wellbeing, as does continuing one's professional life. When factors that might influence quality of life were analyzed, it was found that time after discharge had a positive effect on wellbeing.

Conclusions: Our study revealed that the level of quality of life was low, especially in physical status, and that loneliness was a diminishing factor.

Keywords: Patient; Quality of life; Tracheostomy

Table 4: Effect of descriptive characteristics on mean QOL score.

Variables	Common well-being		Functional status		Symptoms	
	X ± SD	p	X ± SD	p	X ± SD	p
Age						
18-39	41.6 ± 0.0		46.6 ± 4.6		39.3 ± 2.3	
40-69	47.6 ± 14.7	0.798	38.12 ± 6.8	0.07	31.9 ± 5.7	0.048
>70	46.6 ± 5.8		40.0 ± 5.8		30.7 ± 4.9	
Educational status						
No graduation	33.3 ± 0.0		41.0 ± 0.0		33.0 ± 0.0	
Primary school	48.2 ± 15.3	0.094	39.0 ± 6.6	0.85	32.0 ± 6.1	0.863
Secondary and higher education	50.0 ± 8.3		39.6 ± 9.5		33.6 ± 4.3	
Marital status						
Married	49.6 ± 15.5	0.186	38.7 ± 7.2	0.799	31.6 ± 5.7	0.374
Single/Divorced/Widow	42.4 ± 9.9		40.5 ± 5.5		33.9 ± 5.8	
Employment status						
Employed	52.0 ± 7.9	0.184	34.5 ± 3.0	0.046	32.5 ± 4.1	0.879
Unemployed/retired	47.1 ± 15.0		39.8 ± 6.8		32.2 ± 5.9	
Have child						
No children	45.0 ± 8.9		39.9 ± 5.3		30.9 ± 4.6	
1-2 children	47.6 ± 17.7	0.734	38.1 ± 8.7	0.875	34.0 ± 7.2	0.072
>3 children	50.0 ± 14.6		40.0 ± 5.3		31.1 ± 4.2	
Living arrangement						
Living alone	38.0 ± 6.5	0.018	42.7 ± 5.1	0.192	37.0 ± 5.6	0.009
With family	50.0 ± 14.9		38.4 ± 6.8		31.1 ± 5.6	
Location of residence						
Village	33.3 ± 0.0		38.0 ± 0.0		35.0 ± 0.0	
Town	49.0 ± 11.3	0.047	40.2 ± 5.4	0.422	32.3 ± 4.9	0.556
City	46.8 ± 22.6		36.3 ± 10.3		31.6 ± 8.5	

Table 5: Effect of variables on QOL mean score.

Variables	Common well-being		Functional status		Symptoms	
	X ± SD	P	X ± SD	p	X ± SD	P
Own Knowledge about disease						
Enough	47.3 ± 15.2		37.3 ± 6.1		31.2 ± 4.8	
Partially enough	48.1 ± 16.0	0.84	42.3 ± 6.8	0.167	33.8 ± 6.9	0.305
Not enough	48.3 ± 9.1		42.0 ± 7.1		34.0 ± 7.4	
Time after discharge						
0-3 months	50.0 ± 8.0		39.5 ± 5.7		31.9 ± 4.2	
4-8 months	35.7 ± 6.3	0.003	38.4 ± 2.6	0.604	35.5 ± 3.9	0.207
>9 months	51.5 ± 21.6		39.3 ± 9.9		30.8 ± 8.0	
Other health problems						
Have	44.2 ± 9.4	0.348	39.6 ± 4.9	0.835	33.0 ± 4.7	0.823
Have not	50.4 ± 17.2		38.9 ± 8.0		31.7 ± 6.4	
Feeling need for psychological support						
Feel	47.5 ± 12.9	0.948	39.5 ± 6.8	0.631	32.1 ± 5.3	0.823
Doesn't feel	47.9 ± 16.5		38.8 ± 6.8		32.5 ± 6.3	

Table 1: Descriptive characteristics of participants.

Variables	Number (N=36)	Percent (%)
Gender		
Female	3	8.3
Male	33	91.7
Age		
18-39	3	8.3
40-69	25	69.4
>70	8	22.2
Educational status		
No graduation	2	5.6
Primary school	29	80.6
Secondary and higher education	5	13.9
Marital status		
Married	26	72.2
Single/widow	10	27.8
Having children		
No children	10	27.8
1-2 children	14	38.9
3 children	12	33.3
Employment status		
Employed	4	11.1
Unemployed/retired	32	88.9
Living arrangement		
Living alone	7	19.4
With family	29	80.6
Location of residence		
Village	2	5.6
Town	26	72.2
City	8	22.2
Total	36	100

Table 2: Factors that affect QOL.

Variables	Number (N=36)	Percent (%)
Own knowledge about disease		
Enough	22	61.1
Partially enough	9	25
Not enough	5	13.9
Time after discharge		
0-3 months	18	50
4-8 months	7	19.4
>9 months	11	30.6
Presence of other health problems		
Yes	16	44.4
No	20	55.6
Feeling need for psychological support		
Feels	20	55.6
Doesn't feel	16	44.4
Total	36	100

Table 3: Average QOL score of patients.

QOL Subscales	X	SD	Min	Max
Common well-being	47.6	14.4	0	100
Functional status				
Physical function	36.1	19.9	0	100
Role performance	40.7	20.8	0	100
Emotional status	48.1	24.4	0	100
Cognitive status	74	13.4	0	100
Social conditions	47.2	23.4	0	100
Symptoms				
Fatigue	70.3	20	0	100
Nausea and vomiting	22.6	23.9	0	100
Pain	52.3	23.9	0	100
Dyspnea	60.1	31.6	0	100
Insomnia	69.4	24.3	0	100
Anorexia	51.8	26.9	0	100
Constipation	19.4	21.6	0	100
Diarrhea	12.9	27.9	0	100
Economic difficulties	62.9	30.6	0	100

Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duesz NJ, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality of life instrument for use in international clinical trials in oncology. Journal of the National Cancer Institute. 1993; 85:365-76.

Beser NG, Oz F. Anxiety-depression levels and quality of life patients with lymphoma who are curing chemotherapy. J Cumhuriyet Univ Sch Nurs. 2003; 7:47-58.