A study of Otolaryngology emergency admission workload at a busy London hospital

W Jia, A Mohammed, O Olatokun, T Tatla

Introduction

Otolaryngology emergencies are common. Most patients presenting as an emergency do not require admission into the hospital (1)

Objectives

This is an epidemiological study of the patients requiring admission by the Otolaryngology department at Northwick Park Hospital.

Methods

This retrospective study reviewed patient records over a 13 months period (Sept 2016 – Sept 2017), obtained from the hospital coding department.

Results

1370 patients were studied. 53.9% and 46.1% were male and female patients respectively. 84.74% and 15.26% were adult and child admissions respectively. The average length of stay was 2.24 days. Figure 1 shows the overall data. Figure 2 illustrates the monthly distribution of admissions over thirteen months. Table 1 and graph 3 demonstrates the prevalence of specific conditions admitted into the hospital. Upper airway inflammatory conditions, epistaxis and ear related infections are responsible for 67.89% of inpatient admissions.

Conclusions

The vast majority of admissions are relatively benign in nature and require no surgical intervention. Health informatics can help to plan future services.

Figure 1: 866 (63.3%) laryngology (H&N), 358 (26.17%) rhinology and 144 (10.53%) otology admissions from September 2016 to September 2017.

Figure 2: Distribution of patients admitted per month from September 2016 to September 2017. Two peaks of increased admissions were noted in March and July.

Condition	Number of patients	Overall proportion of otolaryngology admissions
Inflammatory / infective upper airway conditions	540	39.42%
Epistaxis	268	19.56%
Otology infections	122	8.91%
Post Tonsillectomy bleeding	104	7.59%
Foreign body in throat	82	5.99%
Neck infections	60	4.38%
Midface infection	43	3.14%
Laryngology (H&N) miscellaneous conditions	39	2.85%
Foreign body in nose	24	1.75%
Airway difficulties	22	1.61%
Otology miscellaneous	22	1.61%
Midface trauma	12	0.88%
Rhinology miscellaneous	11	0.80%
Post Tonsillectomy pain	11	0.80%
Dysphagia	10	0.73%

Table 1: Breakdown of the prevalence of emergency in-patient ENT admissions from September 2016 to September 2017.

Figure 3: Blue bars showing the proportion of different emergency ENT admissions. Red bars showing the proportion of patients requiring a surgical intervention under general anaesthesia.

References

1. Rourke T¹, Tassone P, Philpott C, Bath A. ENT cases seen at a local 'walk-in centre': a one year review. J Laryngol Otol. 2009 Mar;123(3):339-42, doi: 10.1017/S0022215108002508.