Cricothyroidotomy incision length in female subjects - Are the Difficult Airway Society guidelines justified?

P. Fennessy¹, C. Mc Caul¹
¹Mater Misericordiae University Hospital - Dublin (Ireland)

Background and Goal of Study

An 8-10cm incision is recommended for vertical incision cricothyroidotomy when anatomy is indistinct as per the Difficult Airway Society Guidelines 2015. These guidelines are based on expert opinion. At present no evidence-based guideline exists regarding optimum site or length. We investigated clinician accuracy in identification of the cricothyroid membrane (CTM) of female patients.

Failed intubation, failed oxygenation in the paralysed, anaesthetised patient

CALL FOR HELP

Continue 100% O₂
Declare CICO

Plan D: Emergency front of neck access

Continue to give oxygen via upper airway
Ensure neuromuscular blockade
Position patient to extend neck

Scalpel cricothyroidotomy

Equipment: 1. Scalpel (number 10 blade)
2. Bougie
3. Tube (cuffed 6.0mm ID)

Laryngeal handshake to identify cricothyroid membrane

Palpable cricothyroid membrane
Transverse stab incision through cricothyroid membrane
Turn blade through 90° (sharp edge caudally)
Slide coude tip of bougie along blade into trachea
Railroad lubricated 6.0mm cuffed tracheal tube into trachea
Ventilate, inflate cuff and confirm position with capnography
Secure tube

Impalpable cricothyroid membrane
Make an 8-10cm vertical skin incision, caudad to cephalad
Use blunt dissection with fingers of both hands to separate tissues
Identify and stabilise the larynx
Proceed with technique for palpable cricothyroid membrane as above

Materials and Methods

We asked clinicians to identify the CTM, in both the neutral and extended head positions. The distance between the actual (measured using ultrasound) and estimated location of the CTM was measured for both patient positions. We assessed ‘capture’ of the CTM within theoretical incisions of 0-10cm in length using the best clinician estimate as the midpoint around which the incision would be performed. Our aim was to determine the optimum length of incision to capture all measured CTM locations.

Results and Discussion

90 patients were recruited for this study. 66 patients were assessed in both the neutral and extended head position. A further 24 patients had 36 assessments in the neutral head position. In total 168 CTM assessments were performed. The CTM was correctly identified in 18/102 (17.6%) neutral position assessments and 14/66 (21.2%) of the extended position assessments. Clinician inaccuracy ranged from 3.2cm above and 4cm below the actual CTM location across both patient groups.

	Neutral	Extended
Total Assessments N	102	66
All Accuracy N (%)	18/102 (17.6)	14/66 (21.2)
Consultant Assessments N (%)	22/102 (21.6)	11/66 (16.7)
Consultant Accuracy N (%)	9/22 (40.9)	3/11 (27.3)
Senior Trainee Assessments N (%)	63/102 (61.8)	49/66 (74.2)
Senior Trainee Accuracy N (%)	7/63 (11.1)	9/49 (18.4)
Junior Trainee Assessments N (%)	17/102 (16.6)	6/66 (9.1)
Junior Trainee Accuracy N (%)	2/17 (11.8)	2/6 (33.3)
Obese Accuracy N (%)	3/23 (13.0)	2/19 (10.5)
Non-Obese Accuracy N (%)	15/79 (18.9)	12/47 (25.5)
Above CTM N (%)	36/102 (35.3)	17/66 (25.8)
Below CTM N (%)	47/102 (46.1)	34/66 (51.5)
US-DP Distance (cm)	1.17 (-3.2 to +2.8)	1.19 (-4.0 to +2.5)

Conclusion

Using theoretical incisions of increasing length (1cm intervals), an incision size of 8cm with the best clinician estimate as the midpoint would capture all CTM locations in both neutral and extended head positions. Our data thus supports the DAS guidelines.