The Shock Index as a predictor of massive transfusion in trauma patients

I. Otero¹, M. Koo Gómez¹, A. Bonet Burguera¹, R. Reyes García¹, L. Pérez García¹, L. Pariente Juste¹

¹Hospital Universitari de Bellvitge - Barcelona (Spain)

BACKGROUND

Hemorrhage is the leading cause of shock in trauma patients and thereby the main cause of death. The Shock Index (SI) is defined as the ratio of heart rate to systolic blood pressure and, as it has been established previously, a useful tool for diagnosing hypovolemia. The objective of this study is to establish if the SI measured at the Prehospital scenario (PH-SI), at the Hospital admission (H-SI) and at the Intensive Care Unit (C-SI) is a good predictor of red blood units (RBU) and massive transfusion requirements.

MATERIAL AND METHODS

We conducted a retrospective study including all trauma patients from our data base who arrived to hospital from January 2013 to December 2016, under the activation of a Trauma Code and who required admission into the Intensive Care Unit.

Our Hospital is a level 3 trauma center with fully capability of trauma management, part of the reference hospitals of the Trauma Code, which is a code activated by the prehospital personnel and identifies patients with trauma injury, stratifying the severity of the same and the proper center required for its attendance.

SI was calculated at the prehospital, admittance to hospital and critical care settings. The first 24 hours transfused red blood units were recorded and patients were divided into groups according to its amount (less than 5; 5 to 9; and 10 or more matching the general definition of massive transfusion). SI and RBU transfused in patients who needed so were compared with the analysis of the variance and with the Spearman correlation test. A p-value ≤ 0.05 was considered to be significant.

RESULTS AND DISCUSSION

We recruited 184 patients 81.5% of whom were male and 18.5% female, the median age was 43.5 (31- 59) and the mean Injury Severity Score was 32,27; median 29 (22- 36). Their SI values were as follows: PH-SI: 0.77 (0.61 - 1.01); H-SI: 0.78 (0.64 - 1.00); C-SI: 0.92 (0.76 - 1.13). The mortality rate was 14% measured at the seventh day with a total of 26 patients deceased. Patients who met criteria and were transfused were 75. This transfused patients received a median of 4 (2-6) RBC and were categorized in the 3 groups previously mentioned. The Spearman correlation index comparing RBU according to the value of the PH-IS, H-IS and C-IS were 0.46 ($p < 0.001$), 0.66 ($p < 0.001$) and 0.295 ($p = 0.02$) respectively.

RBU	<5	5-<10	≥ 10	p
	n=41	n=21	n=13	
PH- SI	0.86 (0.69 - 1.10)	1.20 (0.78 - 1.32)	1.33 (0.61 - 1.76)	0.013
H- SI	0.91 (0.74 - 1.22)	1.16 (0.94 - 1.57)	1.32 (0.94 - 1.81)	0.009
C-SI	0.97 (0.83 - 1.27)	1.24 (1.00 - 1.48)	1.63 (1.08 - 1.64)	0.187

CONCLUSIONS

The PH-SI and H-SI but not the C-SI are good predictors of transfusion requirements in critical trauma patients who received RBU transfusion, the p value was 0,01 for the PH-SI and 0,009 for the H-SI.

Gender	Male	81,5 %
	Female	18,5 %
Age	43,5 (31-59)	
ISS	29 (22-26)	
Mortality	14% (26)	
PH-SI	0,77 (0,61- 1,01)	
H-SI	0,78 (0,64-1,00)	
C-SI	0,92 (0,76 -1,13)	

1. Nunez TC et al. Early prediction of massive transfusion in trauma: simple as ABC (assessment of blood consumption)? J Trauma. 2009; 66: 346 –352. doi: 10.1097/TA.0b013e3181961c35
2. Olausson A, Peterson E, Mitra B, Jennings P, O'Reilly G, Fitzgerald M. Massive transfusion prediction with inclusion of the pre-hospital Shock Index. Injury. 2015 May;46(5):822-6