Impact of Optimum Diabetes Care on Safety of Fasting Ramadan in Patients with Coronary Heart Disease

Mohamed Hassanein, Fauzia Rashid, Mohammed Abdellatif, Alaaeldin Basheir, Maryam Al Saeed,
Elamin Abdelgadir, Sona Abulkheir, Azza Khalifa, Fawzi Altayb, Fatima AlSayyah, Susan Jacob , Fatheya Alawadi

Dubai Hospital, Endocrinology Department, Dubai, United Arab Emirates

Background

Most of Muslims patients with diabetes and Coronary Heart Disease(CHD) elect to fast in Ramadan, but the actual risk in the different subset of patients with CHD is largely unknown. Theoretically, the extended fasting hours during Ramadan may increase the possibility of hyperviscosity and thrombosis. This factor along with possible missed or altered timings for essential medicines and occurrence of hypoglycemia can result in adverse cardiovascular effects. Only a few studies are done in patients with diabetes and Coronary heart disease (CHD) with an aim to observe any adverse cardiovascular outcome in Ramadan (2,3,4). To the best of our knowledge, no study is done so far to monitor the effect of Ramadan fasting on glycemic control , frequency and severity of hypoglycemia in these high-risk patients with the use of continuous glucose monitor(CGM) and to correlate it with an adverse cardiovascular event in Ramadan fasting.

Aim

We sought to understand the safety of fasting in CHD patients with diabetes insisting on fasting Ramadan. We also monitored the change in biophysical and biochemical parameters of these patients before and after Ramadan.

We conducted this prospective study in a tertiary care hospital in Dubai during Ramadan 2016 that from June 6th till July 5th.

Method

Patient with T2DM with stable known CHD since last three months and insisted on fasting despite advice against it. They were randomly selected from endocrine and cardiology clinic 1 to 2 months before Ramadan, counselled, and asked to sign an informed consent. Patients with the concurrent renal disease or any recent hospitalisation in last 3 months were excluded. All the patients received a session for Ramadan-focused education. The Libre flash sensor insertion is done for continuous glucose monitoring (FSL-CGM) . Their biophysical and biochemical profile was collected 2 to 4 weeks before and after Ramadan.

We recorded DM or CVD-related emergency visit or hospitalisation, change in BMI, systolic and diastolic BP, HBA1c, and frequency of hypoglycemia during Ramadan fasting and not -fasting period. All the quantitative values are described as mean, and analysis of CGM data is shown in frequencies.

Results and Discussion

We recruited 26 patients. There were 21 final participants, 15 males and 6 females with mean age of 58 years (range 34-71). An average number of days fasting were 28. 90% patients (n=18) completed >20 days fasting. Most patients had 3 sensors attached to them with average 3 weeks during Ramadan and 3 weeks outside Ramadan.

Parameter	Pre Ramadan	Post Ramadan
Cholesterol(mg/dl)	140	153
LDL(mg/dl)	73	77
Triglycerides (mg/dl)	130	145
HDL(mg/dl)	42	47
Creatinine(mg/dl)	0.81	0.9
e-GFR	94	91
UREA(mg/dl)	25	28
HBA1c %	7.9	7.3
LFT	N	N

Total 67 episodes of hypoglycemia were recorded during Ramadan. While during non-Ramadan period 24 hypoglycemic events were traced. The average duration for hypoglycemia was 118 minutes (30-340) in Ramadan and 49 minutes (30-184) in non-Ramadan. Further sub-analysis of 8 patients who were found to have 5 or more episodes of hypoglycemia during Ramadan showed, 3 of them were on intense insulin therapy, 3 were receiving SU based oral therapy and 2 on basal insulin in combination with metformin and DPP4 inhibitors. But those on SU based therapy never had any hypoglycemia outside Ramadan unlike to the patients on insulin based regimen who did have hypoglycemic episodes during non Ramadan period as well, though relatively fewer in number. None of them reported severe hypoglycemia, aggravation of their underlying coronary heart disease or chest pain during the Ramadan fasting, only one patient visited emergency department with dyspnea after Ramadan.

Frequency of Hypoglycemia

Hypoglycemia Timing

Conclusion

This is first study ever done in high-risk CHD patients to objectively assess the glycemic changes, frequency of hypoglycemia and its correlation with any adverse CVD effect through the use of FSL-CGM . Patients had 2.7 times higher incidence of hypoglycemia that is more marked and prolonged during fasting. Those on sulfonylurea bases treatment demonstrated more severe and frequent hypoglycemia during Ramadan compared to non -Ramadan. No significant alteration was seen in BMI, SBP and DBP, HBA1c, lipid profile and hepatic functions. Studies with larger sample size needed for further validation of these findings.

References

1. International Diabetes Federation. Diabetes and cardiovascular disease. Brussels, Belgium: International Diabetes Federation, 2016. www.idf.org/cvd
2. Chamsi-Pasha M, Chamsi-Pasha H. The cardiac patient in Ramadan. *Avicenna J Med*. 2016;6(2):33.
3. Mousavi M, Mirkarimi S, Rahmani G, Hosseinzadeh E, Salahi N. Ramadan fast in patients with coronary artery disease. *Iran Red Crescent Med J*. 2014 Dec;16(12):e7887.
4. Salim I, Al Suwaidi J, Ghadban W, Alkilani H, Salam AM. Impact of religious Ramadan fasting on cardiovascular disease: a systematic review of the literature. *Curr Med Res Opin*. 2013 Apr;29(4):343-54