The study of the immune profile dynamics and the level of endogenous intoxication in patients with polytrauma

VOLKOVA Yu.V.

Doctor of Medicine, Professor of the Department of Emergency Medicine, Anesthesiology and Intensive Care

BARANOVA N.V.

Candidate of Medical Sciences, assistant of the Department of Emergency Medicine, Anesthesiology and Intensive Care

LANTUHOVA N.D.

Candidate of Medical Sciences, assistant of the Department of Emergency Medicine, Anesthesiology and Intensive Care

KUDINOVA O.V.

Candidate of Medical Sciences, Associate Professor, Department of Emergency Medicine, Anesthesiology and Intensive Care

Kharkiv National Medical University

Kharkiv, Ukraine

Background

Formation of the reaction of the immune system to the polytrauma begins with the first minutes of its receipt, the implementation of adaptive reactions to primary damage with a high risk of multiple organ disfunction development.

The aim of the study:

Study of the immune status level and the severity of the inflammatory response in patients with polytrauma. Measurement of the level of C-reactive protein was carried out by a "sandwich - variant" of solid-phase immuno enzyme analysis.

Methods

The immune status was studied in 95 patients with polytrauma during the first day after receiving an injury(LII, the CD3, CD4, CD8 population, the immunoregulatory index, IgG, IgM according to standard techniques). Measurement of the level of C-reactive protein was carried out by a "sandwich - variant" of solid-phase immuno enzyme analysis.

Results

The index of LII (leukocyte index of intoxication) was studied, that was at the level of 0.91 ± 0.08 during the first day (norm - 0.8-1.6). Increasing of the C-reactive protein in the level of 8 times and amounted to 24.6 ± 4.1 mg/l (norm-0-3 mg/l). The analysis of primary cellular indicators and humoral immunity revealed that the relative number of T lymphocytes CD3 was $53.4 \pm 4.2\%$, which was 21.5% lower than the norm of 58-76%.

Expression of the populations of helpers and suppressors was initially reduced by 11.8% and amounted to $20.3 \pm 2.6\%$, for the CD4 population (norm-23-48%), decreased by 6.5% - $15.9 \pm 1.6\%$ for the CD8 population respectively(norm-17-25%). The decrease in immunoreactivity was confirmed by low values of the immunoregulatory index - 1.28 ± 0.02 (norm-1.8-2.5). The level of immunoglobulin fractions was within the normal range: for IgM - 1.01 ± 0.03 g/l(norm-1,19-2,4 g/l). For IgG values were 6.6 ± 0.3 g/l(norm -8-17g/l).

Conclusion

Thus, in response to traumatic effects, on the one hand, there was a pronounced inflammatory stimulation characterized by high C-reactive protein that indicates the initiation of an inflammatory response within the next hours after the injury, on the other hand, there was the hyporeactivity of the cellular and humoral responses.