Pseudocontinuous arterial spin labeling on magnetic resonance imaging in patients with carotid stenosis

SF Cheng, A Di Napoli, T Richards, MM Brown, HR Jäger

Division of Surgery and Interventional Science & Institute of Neurology, University College London, United Kingdom
Neurosciences, Mental Health and Sensory Organs Department, Sapienza University, Rome, Italy

Background

- Arterial transit artefacts (ATA) are high signals on the surface of the brain, which is labeled blood that has not reached the capillary bed (gives an indication of brain perfusion).
- Aim: To determine the presence of ATAs on MRI in symptomatic and asymptomatic carotid stenosis and to evaluate its significance in these patients.

Methods

- 45 patients with internal carotid stenosis $\geq 50\%$ who had high resolution MRI perfusion imaging done were retrospectively included.
- Patients were divided into symptomatic and asymptomatic carotid stenosis, 50-70% and $\geq 70\%$ stenosis.
- PcASL images were analysed and scored on the presence of ATA's in both hemispheres.
- The circle of Willis was scored on completeness (0: no Acom or Pcom; 1: Acom or Pcom; 2: both present).

Figure 1. Pseudocontinuous arterial spin labelling on MRI.

		ATA absent (n=29, %)	ATA present (n=16, %)	P- value
Age (years)		67.9 $\pm$ 9.0	76.2 $\pm$ 8.0	0.0038
Male gender		20 (69)	12/16 (75)	0.7432
Ulcerated/irregular plaque		9 (31)	8/16 (50)	0.3356
Stenosis >70%		11 (38)	16/16 (100)	<0.000 1
Intra plaque haemorrhage		8 (28)	4/16 (25)	1
Circle of Willis	0	2 (7)	9/16 (56)	0.0005
	1	19 (68)	7/16 (44)	
	2	7 (25)	0/16 (0)	
PCOM (contralateral)		6 (21)	3/16 (19)	1
PCOM (side of interest)		10 (36)	2/16 (13)	0.1601
ACOM		23 (82)	5/16 (31)	0.0012
Symptomatic		10 (34%)	13 (81%)	0.005

Table 1. Patients with and without ATAs on MRI.

Arterial Transit Artefact scoring system	
0	No or minimal ASL signal
1	Moderate ASL signal with ATA
2	High ASL signal with ATA
3	Normal perfusion without ATA

Figure 2. Arterial Transit Artefact Scoring System

Results

- Patients with $\geq 70\%$ carotid stenosis all had ATAs (n=16) compared to 38% of the patients with 50-70% stenosis (n=11).
- Symptomatic patients were associated with the presence of ATAs (81% in symptomatic vs. 34% asymptomatic; p=0.005)

Conclusion

Only patients $\geq 70\%$ carotid stenosis were shown to have ATA's. Patients with symptomatic carotid stenosis more often had ATAs.

University College London Hospitals

NHS Foundation Trust