Ethnicity and survival from out-of-hospital cardiac arrest (OHCA): A New Zealand registry study.

Graham Howie, Kate Conaglen, Bridget Dicker.

All authors: St John Ambulance, New Zealand, and Auckland University of Technology (AUT), New Zealand.

Objective: To determine the relationships between survival from all-cause out-of-hospital cardiac arrest (OHCA) and ethnicity in New Zealand.

Methods: A retrospective observational study was conducted using data compliant with the Utstein guidelines from the St John Ambulance New Zealand OHCA Registry for adult patients who were treated for an OHCA between 1 October 2013 and 30 September 2015.

Primary outcomes measured were: Return of spontaneous circulation (ROSC) sustained to handover at hospital, and survival to thirty-days, for New Zealanders of these ethnicities: European, indigenous Māori, Pasifika (Pacific Islanders), and Other. Standard Utstein elements were also probed, such as: locality of arrest, whether or not the OHCA was witnessed, CPR and defibrillation prior to ambulance arrival, initial rhythm, aetiology.

Results:

	New Zealand Population (2013) ¹ <i>n</i> = 4.24 million (%)	OHCA Incidence ³ <i>n</i> = 3,214 (%)	ROSC (%)	30-day Survival (%)
European	74.6	59.5	32.3 ^a	18.1 ^a
Māori	15.6	24.2	26.9 ^b	10.5 ^b
Pasifika	7.8	9.0	28.5 ^{a,b}	13.5 ^{a,b}
Other	14.2	6.9	32.9 ^{a,b}	12.7 ^b
	112.2% ²	100%	<i>p</i> < 0.03	<i>p</i> < 0.001

Same superscript letters denote no significant difference from each other at the 0.5 level.

Pearson Chi-Square test.

Locality: No difference of incidence of arrest in public places. Median time until ambulance arrival (7-8 minutes) was similar for all. Māori and Pasifika were more likely to arrest at home, whereas European were more likely to arrest in aged-care facilities.

Aetiology: Māori were relatively more likely to suffer non-cardiac arrests than either European or Pasifika.

? Witnessed: Māori were more likely to suffer unwitnessed arrest than European or Pasifika. Europeans were more likely to have a witnessed arrest in the presence of ambulance crews, than Māori or Pasifika.

Bystander CPR: Māori and Pasifika were more likely to have CPR prior to ambulance arrival than Europeans.

Initial Rhythm: Europeans were more likely to have an initial shockable rhythm than Māori, but not Pasifika. (All differences: *p* < 0.05 Pearson Chi-Square.)

Conclusions: Utstein-favourable factors such as initial shockable rhythm, witnessed arrest and bystander-CPR may contribute to these differences in survival from OHCA between ethnicities. Proposed additional factors: socioeconomic status⁴, cardiovascular risk factors^{5,6}, and rural/urban localities^{7,8}.

References: 1. <http://archive.stats.govt.nz/~media/Statistics/browse-categories/snapshots-of-nz/nz-in-profile/2015/nzip-2015.pdf> 2. <http://archive.stats.govt.nz/~media/Statistics/surveys-and-methods/methods/class-stnd/ethnicity/ETHNIC05-statistical-standard%20v2%20cls.pdf>

3. See <https://www.stjohn.org.nz/News-Info/Our-Performance/Cardiac-Arrest-Annual-Report/> for annual St John Ambulance New Zealand OHCA Registry Reports 4. Merchant RM, Becker LB, Yang F, Groeneveld PW. (2011). Hospital racial composition: A neglected factor in cardiac arrest survival disparities. *Am Heart J* 2011;161(4):705-711. doi: org/10.1016/j.ahj.2011.01.011

5. Bullen C, Beaglehole R. Ethnic differences in coronary heart disease case fatality rates in Auckland. *Aust NZ J Pub Health* 1997;21(7):688-693. doi:10.1111/j.1467-842X.1997.tb01781.x

6. Curtis E, Harwood M, Riddell T. et al. Access and society as determinants of ischaemic heart disease in indigenous populations. *Heart, Lung and Circulation* 2010;19(5):316-324. doi: org/10.1016/j.hlc.2010.04.129

7. Masterson S, Wright P, O'Donnell C. et al. Urban and rural differences in out-of-hospital cardiac arrest in Ireland. *Resuscitation* 2015;91:42-7. doi:10.1016/j.resuscitation.2015.03.012

8. Mathiesen WT, Bjørshol CA, Kvaløy JT, Søreide E. Effects of modifiable prehospital factors on survival after out-of-hospital cardiac arrest in rural versus urban areas. *Critical Care* 2018;22(1):99. doi:10.1186/s13054-018-2017-x.