Chelation of the toxic aluminum (Al^{3+}) by Clofibric acid (CA) in Aqueous media

Y. Z. Hamada¹, A. Ayesh¹, and M. Z. Badr² ¹LeMoyne-Owen College, Memphis, TN 38126 and ²Division of Pharmacology & Toxicology, School of Pharmacy, University of Missouri-Kansas City, Kansas City, MO 64108

Chemistry

LeMoyne-Owen
COLLEGE

Abstract

This presentation is part of the Peroxisome Proliferator-Activated Receptors- α (PPAR- α) research. PPARs are a group of receptor proteins that function as transcription factors regulating the expression of genes. From our on-going efforts to study the low molecular mass ligands that binds to these PPARs, we are presenting the reactions of CA with Al^{3+} in aqueous solutions. The objectives of this work are: 1- to find whether there is a binding between Al^{3+} and CA, and 2- to find what is the binding mode if there is binding. After using the potentiometric technique, the UV-Vis, and IR spectroscopies it appeared that there is strong binding at which a series of ternary Al-CA-OH species are formed. The main complex was $[\text{Al}(\text{CA})(\text{OH})_3]^-$. These data will be discussed in details.

Experimental

The potentiometric titration solutions were contained in a 250 mL beaker equipped with a magnetic stirring bar. The beaker was covered with a custom made Teflon cover. In a typical titration; Clofibric acid ligand (CA) was added first followed by the Al^{3+} solution. D. I water was added to take the total volume to 100 mL. Details of solution preparation, standardizations and potentiometric titrations have been documented in references 1-4. All IR spectroscopy measurements were conducted using Nicolet iS10 spectrophotometer in connection with OMNIC software version 8.1, both purchased from Thermo Fisher Scientific (Madison, WI). The UV-Vis experiments were conducted on Thermo Scientific *Evolution array spectrophotometer* using 4.0 mL set of Quartz Cuvettes.

Al^{3+} /CA mole ratio	Number of titration runs	Vol. (mL) of NaOH titrant (0.1062 M)	Equivalents of NaOH titrant	Proposed species	Remarks
1:0	4	1.62	3.02	$\text{Al}(\text{OH})_3$	Three H^+ were released
0:1	4	0.62	1.04	$[\text{CA}-\text{COO}^-]$	Clofibric acid is a mono-protic acid
1:1	3	3.30	3.50	$[\text{AlCA}(\text{OH})_2]$	The ternary (Al-CA-OH) complex is formed
1:2	6	3.50	4.00	$[\text{AlCA}(\text{OH})_2]$	The ternary (Al-CA-OH) complex is formed or a dimmer
1:3	4	4.04	4.70	$[\text{AlCA}(\text{OH})_2]$	The ternary (Al-CA-OH) complex is formed
1:4	4	4.5	5.00	$[\text{AlCA}(\text{OH})_2]$	The ternary (Al-CA-OH) complex is formed

References

- [1] Hamada, Y.Z.; Harris, W.R.; Rath, N. Crystal Structure of Pyridoxal Amino Methyl Phosphonic Acid (PYRAMPA) and Its Stability Constants with Al^{3+} . *International J. Green and Nano Tech.* (1) 1-8 **2013**.
- [2] Hamada, Y.Z. et al, Equilibrium models of Cr^{3+} and Cu^{2+} with glutamate. *J. Coord. Chem.*, Vol. 62 (5) 721-733 **2009**.
- [3] Hamada, Y.Z. and Rogers, C. *J. Coord. Chem.*, Vol. 60 (20) 2149-2163 **2007**.
- [4] Hamada, Y.Z. et al., Accurate potentiometric studies of chromium-citrate and ferric-citrate complexes in aqueous solutions at physiological and alkaline pH-values , *Synthesis and reactivity of inorganic and metal-organic and nano-metal chemistry* Vol. 36, 469-476 **2006**.
- [5] Harris, W.R. and Hamada, Y.Z. Stability Constants of Aluminum with Phosphonic Acid Derivatives and Multinuclear NMR-Measurements in Aqueous Solutions”, *Inorg. Chim. Acta.* Vol. 359 (4) 1135-1146 **2006**.
- [6] Hamada, Y.Z. S. Rehan, J. Scott Clofibric Acid, a Peroxisome Proliferator-Activated Receptor α Agonist, forms a Ternary Complex with Ferric Iron Implications for the Cardiovascular Protective Effect and Toxicity of Fibrates, *Electronic Journal of Biology (eJBio)*, S2:18-24 **2016**.
- [7] Hamada, Y.Z.; Badr, M.Z; Hayes, J.; Yates, B. Ternary Metal-Hydroxo Chelate of Cr^{3+} with Clofibric acid (CA): a Peroxisome proliferator-activated receptors- α (PPAR α) ligand. *J Heavy. Met. Tox. Disease*, Vol. 1 No1:6, 1-8 **2016**.

Acknowledgements

NSF # HRD-1332459

ACS-SEED Program

Conclusion

- [1] Calibrating the potentiometric titration system with H_3PO_4 proved system accuracy.
- [2] Free Clofibric Acid has generated 1.0 proton out of the single carboxylic acid group.
- [3] 3.5 protons appeared from the 1:1 runs and 4.0 protons appeared from the 1:2 runs.
- [4] Both the potentiometric titrations and the IR indicated that the complex formed are not $\text{Al}(\text{OH})_3$. This is very consistent with previous data collected [5-7].
- [5] Determining the exact structures for these new complexes is our future work.