The role of DHN-melanin and RodA in the protection of *Aspergillus fumigatus* conidia against Reactive Oxygen Species

E. M. Keizer, I.D. Valdes, M. Schinkel, F. van Baelen, H. Wösten, H. de Cock

Introduction: *Aspergillus fumigatus* conidia entering and invading lung epithelium use SOD and catalases to defend against ROS. DHN-melanin (Fig 1A) was previously reported to protect against ROS¹ but the mechanism remained unclear. Interestingly, L-DOPA melanin (Fig. 1B) produced by *Cryptococcus neoformans* protects against hydroxyl radicals by acting as an antioxidant² (Fig. 1C). Here we investigated ROS sensitivity of a $\Delta pksP::hph$ mutant lacking DHN-melanin and compared it to a *pksP* mutant obtained via UV mutagenesis.

Fig 1A. Biosynthesis of DHN-melanin

Fig 1B. Biosynthesis of L-DOPA melanin

Fig 1C. Oxidation states of quinonic residues in L-DOPA melanin³

$\Delta pksP::hph$ or $\Delta rodA::hph$ mutants have no increased sensitivity to H_2O_2

Left: Growth inhibition zones of strains due to 500 mM hydrogen peroxide (in center hole). **Right:** Average diameter (mm) of the observed inhibition zones. Note that the UV *pksP* mutant is more sensitive like previously reported¹.

No role for *A. fumigatus* conidial surface components DHN-melanin or RodA in protection against hydroxyl radicals

Conclusions

In contrast to L-DOPA melanin, DHN melanin does not protect conidia of *A. fumigatus* conidia against H_2O_2 or hydroxyl radicals. Differences in ROS sensitivity between *pksP::hph* and the UV mutant are likely due to other mutations.