Influence of bacterial resistance on mortality in intensive care units: A registry study from 2000 to 2013 (IICU Study)

Bonnet V.¹, Pasqualini L.¹, Dupont H.², Kipnis E.³, Hanouz J.-L.¹, Fischer M.-O.¹

1University Hospital Caen, Dept of Anaesthesiology & Intensive Care, Caen, France, 2University Hospital Amiens, Dept of Anaesthesiology & Intensive Care, Amiens, France, 3University Hospital Lille, Dept of Anaesthesiology & Intensive Care, Lille, France

Background and Goal of Study: Bacterial resistance to antibiotics is a daily concern in intensive care units and represent a major public health challenge in every country, with the emergence of pan-drug resistant bacteria. However, few data are available concerning clinical consequences of in-vitro defined resistances. This wide retrospective study aimed at comparing in-hospital mortality of intensive care infections, according to bacterial resistance profile.

Materials and Methods: Health-care associated infections are monitored in a national prospective registry, managed by the Investigation and Surveillance of Healthcare-Associated Infections Network (RAISIN). Data included in the registries from 2000 to 2013 were analysed. Sub-group analysis was performed using one-way ANOVA.

Pathogen	Incidence (%)	95% CI	Resistance (%)	95% CI
Staphylococcus aureus	19.9	[19.0 - 20.8]	30.7	[28.5 - 32.9]
Pseudomonas aeruginosa	15.7	[14.9 - 16.5]	43.9	[41.2 - 46.6]
Enterobacter spp.	7.2	[6.6 - 7.8]	72.6	[69.0 - 76.2]
Klebsiella pneumoniae	3.1	[2.7 - 3.5]	64.9	[59.1 - 70.7]
Enterococcus faecalis	2.2	[1.9 - 2.5]	8.6	[4.6 - 12.6]
Acinetobacter baumannii	1.1	[0.9 - 1.3]	70.5	[61.3 - 79.7]
Enterococcus faecium	0.6	[0.4 - 0.8]	49.0	[35.0 - 63.0]

Results and Discussion: Among the 87931 patients retrieved, 10001 presented a health-care related infection, including 3092 resistant germs (36.7% (95%CI [35.7-37.7])). Infections were caused by gram negative bacilli in 54.7% (95%CI [53.6 - 54.8]) of cases and gram positive cocci in 30.0% (95%CI [29.0-31.0]) of cases. Gram negative bacilli presented the highest rate of resistance (52.8% 95%CI [51.3-54.3]), in comparison to Gram positive cocci (48.1% 95%CI [46.3 - 49.9]). In-hospital mortality was higher in case of infection if the germ had antibiotic resistance (51.9% 95%CI [50.1-53.7] vs 45.5% 95%CI [44.2-46.8], p< .001) and critical care length of stay was longer (33.5 ± 26.0 vs 29.5 ± 22.9 days, p< .001). These results remained significant after SAPS II matching (p< .001), and were identical in the gram negative bacilli and gram positive cocci sub-groups. Comparing patients according to their origin before admission, no difference of mortality was found. However, more resistant bacteria were found in patients coming from intensive care units (45.8% 95%CI [41.5-50.1]) or conventional units (39.3% 95%CI [37.6-41]) than from home (33.5% 95%CI [32.1-34.9]) or long term care or rehabilitation centres (33.5% 95%CI [28.8-38.2]). Central venous catheter infections were associated to the highest rate of resistance ((75.0% 95%CI [63.2-86.8]), in comparison to pneumopathies (48.2% 95%CI [46.6-49.8]), bacteraemia (48.7% 95%CI [45.3-52.1]) and urinary tract infections (40.8% 95%CI [37.5-44.1])).

Conclusion: Bacterial resistance is associated with a higher ICU mortality and increased length of stay, regardless of germ or origin of the patient.