Utstein style report and factors influencing the outcome after implementation of the ERC Resuscitation Guidelines 2015

Purpose of the study:

- To estimate the incidence of out-of-hospital cardiac arrest (OHCA) in Ljubljana
- To estimate the outcome of OHCA
- To identify the factors influencing OHCA outcome

Materials and methods:

- Retrospectively recorded data of all patients with OHCA
- Area covered by the Prehospital emergency medical service Ljubljana
- Period 2016-2018
- Statistical methods: descriptive statistics and multivariate logistic regression

Results:

- Cardiopulmonary resuscitation was attempted in 557 patients
- The OHCA was witnessed in 430 cases
- Basic life support was performed in 344 cases
- Defibrillation with an automated external defibrillator before the arrival of the emergency medical service was performed in 53 patients
- 77 patients survived to discharge from hospital
- At the time of discharge 61 patients had cerebral performance category (CPC) 1 or 2

Authors:

- Mateja Škufca Sterle, MD, MSc. Prehospital Emergency Medical Service Ljubljana, Community Health Centre Ljubljana, Slovenia
- Eduard Kralj, MD, PhD. Institute of Forensic Medicine, Faculty of Medicine, University of Ljubljana, Slovenia

Figure:
Utstein standardized template for reporting outcomes from OHCA

Conclusion:

In the studied population, 77 (13.8 %) patients having suffered OHCA survived to discharge from hospital and 61 (10.9 %) had favourable functional outcome. Arrival time of EMS, witnessed arrest, defibrillation before the arrival of EMS and onset of cooling in the field had influence on survival after OHCA.

Reference:

Perkins GD, et al. Cardiac Arrest and Cardiopulmonary Resuscitation Outcome Reports: Update of the Utstein Resuscitation Registry Templates for Out-of-Hospital Cardiac Arrest. Resuscitation 2015;96:328-40.