Impact of perioperative glycemic control target on mortality and major complication after CABG——a study in 10,000 Chinese adults

Yanyan Chen¹ Xiaopei Hou¹ Heng Zhang² Xiaojue Li¹ Xinxing Feng¹ Xin Qian¹ Wei Wang² Hansong Sun² Na Shi¹ Shengshou Hu² Zhe Zheng² Guangwei Li¹

1 Endocrinology and Cardiovascular disease Center, Fuwai Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China;
2 Department of Cardiovascular Surgery Center, Fuwai Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

BACKGROUND AND AIMS

Guidelines recommend a moderate glucose control target of 7.8-10mmol/L for patients with hyperglycemia recovering in the intensive care unit(ICU) after cardiac surgery. It is unclear whether the same target can be used for hyperglycemic patients with and without diabetes.

METHODS

- We studied 9,987 patients including 5,830 patients without diabetes and 4,157 patients with diabetes who underwent coronary artery bypass grafting (CABG) at a single hospital in China from 2011-2014.
- Patients were stratified into three groups according to their mean blood glucose (MBG) levels during the ICU admission: 1) strict-control, MBG <7.8 mmol/L; 2) moderate-control, MBG 7.8-10.0 mmol/L; and 3) liberal-control, MBG ≥10mmol/L.
- The relationship between glycemic control and the composite outcome which including in-hospital mortality and major complication (sum of acute myocardial infarction, stroke and renal failure during the hospitalization after CABG surgery) was investigated by using multivariate logistic regression model adjusting for cardiovascular risk factors.

RESULTS

- Compared with patients without diabetes, patients with diabetes had higher systolic blood pressure, fasting blood glucose at admission, and HbA1c; and more frequently had a history of hypertension, dyslipidemia, myocardial infarction, and three-vessel disease (P<0.01 for all comparisons) (**Table 1**).The rate of composite outcome was 6.1% in the whole population, and occurred more frequently in patients with diabetes than in those without diabetes (7.6% vs 5.0%; P<0.001). The in-hospital mortality and major complication were also more common for patients with diabetes.
- The rate of composite outcome was significantly higher in liberal-control group, while significantly lower in strict-control group, when compared with moderate-control group among patients without diabetes. In diabetes cohort, the rate of composite outcome was significantly higher in both liberal-control and strict-control groups. Among patients without diabetes, in-hospital mortality in liberal-control group was significantly higher than moderate-control group, but patients with diabetes had higher rate in strict-control group. The rate of major complication was consistent with composite outcome in the patients without diabetes, while in the diabetes group, liberal-control group showed significantly higher rate of complication comparing with moderate-control group (**Figure 1**) .
- After adjustment of the multiple risk factors, patients with diabetes in strict-control group had higher risk of in-hospital mortality compared with patients in moderate-control group (odds ratio [OR], 11.04; 95% confidence interval [CI], 4.98-24.46; P<0.001); but among patients without diabetes, liberal-control group had higher risk of death (OR, 7.14; 95% CI, 2.34-21.80; P<0.001). The risk of major complication was significantly lower in strict-control group comparing with moderate-control group in patients without diabetes (OR, 0.72; 95% CI, 0.53 - 0.99; P=0.048), in contrast to patients with diabetes, there was no difference in glycemic control groups (OR, 1.17; 95% CI, 0.68-2.00; P=0.57) (**Figure 2**).

CONCLUSIONS

- Hyperglycemia was independently associated with in-hospital mortality and major complication among CABG patients during the postoperative hospitalization.
- Adverse outcomes were often lower for those with moderate glycemic control (7.8-10.0mmol/L). However, strict glucose control targeted to <7.8mmol/L was associated with increased in-hospital mortality among patients with diabetes, but decreased occurrence of a major complication among patients without diabetes.
- The optimal glycemic control target after CABG may differ by diabetes status.

Table 1. Baseline Characteristics in all patients, patients without and with diabetes

Variable	All Patients (n=9987)	Patients without Diabetes (n=5830)	Patients with Diabetes (n=4157)
Age-y	61 (55-67)	60 (55-66)	61 (55-67)
Male sex	7711/9947(77.5)	4646/5809(79.9)	3065/4138(74.1)
BMI*	25.5 (23.5-27.6)	25.3 (23.3-27.3)	25.8 (23.9-27.9)
SBP mmHg	128 (119-140)	125 (117-140)	130 (120-140)
DBP mmHg	76 (70-80)	76 (70-80)	76 (70-80)
FBG mmol/L	5.5 (4.9-6.6)	5.1 (4.7-5.5)	6.9 (5.8-6.4)
HbA1c %	6.2 (5.8-6.7)	5.9 (5.6-6.1)	7.2 (6.6-8.3)
TG mmol/L	1.5 (1.1-2.0)	1.5 (1.1-2.0)	1.5 (1.1-2.1)
Cholesterol-mmol/L	4.1 (3.5-4.8)	4.1 (3.5-4.9)	4.0 (3.4-4.8)
HDL-c mmol/L	1.0 (0.9-1.2)	1.0 (0.9-1.2)	1.0 (0.8-1.2)
LDL-c mmol/L	2.4 (1.9-3.1)	2.5 (2.0-3.1)	2.4 (1.8-3.0)
Smoker	4263/9950(42.8)	2593/5803(44.7)	17670/4147(40.3)
Hypertension†	6128/9950(61.6)	3402/5803(58.6)	2726/4147(65.7)
Dyslipidemia†	5553/9950(55.8)	2961/5803(51.0)	2592/4147(62.5)
Chronic renal failure†	15/9987(0.15)	8/5830(0.14)	7/4157(0.17)
Previous myocardial infarction†	2631/9950(26.4)	1467/5803(25.3)	1164/4147(28.1)
Previous stroke†	1043/9950(10.5)	543/5803(9.4)	500/4147(12.1)
Peripheral vascular disease†	1209/9950(12.2)	625/5803(10.8)	584/4147(14.1)
Three-vessels disease	2541/9987(25.4)	1386/5830(23.8)	1155/4147(27.8)

*The body-mass index is the weight in kilograms divided by the square of the height in meters.

† Hypertension, dyslipidemia, Chronic renal failure, previous myocardial infarction, previous stroke, previous peripheral vascular disease mean previously diagnosed in other hospitals before admission.

Figure1. Outcomes by glycemic control in the intensive care unit after coronary artery bypass grafting for patients without and with diabetes.

Figure 2. Odds Ratio for composite outcome, in-hospital mortality and major complication in strict-control, moderate-control and liberal-control group.

Multivariable regression analysis for outcomes in all patients, patients without and with diabetes in ICU after adjustment age, sex, smoking, SBP, LDL-c, chronic renal failure, left heart failure, previous myocardial infarction, three-vessel disease, combined valve surgery and extracorporeal circulation. Squares represent odds ratios, and horizontal bars represent 95% confidence intervals. A. liberal-control group compare with moderate-control group. B. strict-control group compared with moderate-control group.

References:

1. Lazar HL, McDonnell ME, Chipkin SR, et al. The Society of Thoracic Surgeons Practice Guideline Series: Blood Glucose Management During Adult Cardiac Surgery[J]. The Annals of Thoracic Surgery. 2009,87(2):663-9.
2. Moghissi ES, Korytkowski MT, Dinardo M, et al. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control[J]. Diabetes Care. 2009,32(6):1119-31.
3. Investigators N-SS, Finfer S, Chittock DR, et al. Intensive versus conventional glucose control in critically ill patients[J]. N Engl J Med. 2009,360(13):1283-97.

Contact details:

Professor Guangwei Li, guangwei_li45@126.com and Professor Yanyan Chen, doc.yan@medmail.com.cn

Funding: Key Program of Discipline Development of Chinese Academy of Medical Sciences