Liraglutide & dulaglutide in addition to SGLT-2i and metformin in Indian T2DM: a retrospective observational study

S. Ghosal¹, B. Sinha².

¹Nightingale Hospital, Endocrinology, Kolkata, India.

²AMRI hospitals, Endocrinology, Kolkata, India.

ABSTRACT:

Aim: The aim of this analysis was to test the additive metabolic benefit of sequential addition of SGLT-2i followed by GLP1-RA along with metformin in a real-life scenario.

Material & Methods: A retrospective real world observational case note study conducted in two diabetes care centres in India analyzed the first 60 consecutive T2D patients who could afford this therapy and had not achieved their glycaemic target on metformin and SGLT2i were added either Dulaglutide or Liraglutide and followed up for 13 weeks.

Results: Both liraglutide and dulaglutide proved to be an excellent add on to metformin and SGLT-2 inhibitor. Liraglutide had an additional significant impact on systolic blood pressure reduction as well as larger proportion achieved target HBA1C in contrast to the dulaglutide.

Conclusion: In this retrospective study in Indian type 2 diabetic patients poorly controlled with metformin and SGLT-2 inhibitor we found a meaningful impact of adding a GLP1-RA on all metabolic parameters. There were additional advantages seen with liraglutide as far achieving target HBA1C of less than 7% and also on the quantum of weight loss and systolic blood pressure reduction.

Keywords:

Diabetes; liraglutide; dulaglutide; SGLT-2i ; HBA1C; weight; blood pressure.

Key Messages:

- I. This is the first real life data on metabolic impact of sequential addition of GLP1-RA after failure of metformin and SGLT-2 inhibitor.
- II. This study demonstrated a significant metabolic benefit of adding either dulaglutide or liraglutide after metformin and SGLT-2 inhibitor failure.
- III. This study also demonstrated superior reduction of weight and systolic blood pressure with liraglutide compared to dulaglutide with larger proportion of patients achieving HBA1C targets in liraglutide arm.
- IV. The prime importance of this study is related to the fact that in India patients have today out of their pocket for medications.

Introduction:

The combination of Glucagon like Peptide Receptor Analogues (GLP-1RA) with a Sodium-glucose linked transporter inhibitor (SGLT2i) addresses six out of the eight ominous pathophysiological defects seen in Type 2 Diabetes (T2D). (1). Combining both these class of drugs seems to address the pathophysiological defects as well address CV risk. (2,3,4,5). Hence there is a good scientific rationale for using them in combination.

The recent DURATION 8 study, testing the above rationale, demonstrated an additive effect on weight and systolic blood pressure reduction but not HBA1C reduction when exenatide LAR and dapagliflozin were used in combination as a co-initiation strategy (6).

In a resource poor setting such as India where patients pay ‘out-of-pocket’, injectable therapies are usually used as a third-line agent when oral therapy fails. Hence, data on this combination mimicking a real-life setting i.e. oral therapy followed by injectable would be useful in guiding physician choices for intensification options in Type 2 diabetes.

Materials & Methods:

A retrospective, real world observational study to evaluate the efficacy of triple-anti-hyperglycemic agent therapy namely metformin, sodium glucose co-transporter 2 Inhibitors (SGLT-2Is) and glucagon like peptide receptor agonists (GLP-1 RAs) for patients failing on a combination of full dose metformin 2000mg/day and SGLT2i, was conducted in the outpatients of two hospitals in Kolkata, India, from May 2016 to August 2016. The baseline characteristics of the patients included for analysis are detailed in table 1 (Supplementary appendix).

After clearance from the local ethics committee, the case notes of the first 30 consecutive patients who had been commenced on Metformin plus SGLT2i plus Dulaglutide plus the first 30 patients who had been commenced on Metformin plus SGLT2i plus Liraglutide was collated.

After adequate counseling patients *who could afford this expensive combination of medications for at least 3 months* had been commenced on this therapy on the basis of the following inclusion and exclusion criteria:

Inclusion Criteria:

Adult Type 2 diabetics with HbA1C≥7.0 % on Metformin plus SGLT2i

BMI ≥ 25 kg/m²,

eGFR>45 ml/min

Exclusion Criteria:

- Type 1 Diabetes
- Pregnancy
- Deranged liver function tests
- Any major organ system disease as determined by physical examination, medical history and screening blood tests
- Recent insulin therapy
- Recent anti obesity therapy
- Recent treatment with any other oral anti diabetics

Patients received treatment as per routine standard of care. All anti hypertensives, anti hyperlipidaemics and anti platelet agents and other preexisting medications were continued as per the patients’ requirements. All patients’ records with respect to age, gender, height, body weight, BMI, duration of diabetes, HbA1c, FPG, BP and adverse effects were collected from the case note database. Blood glucose was measured by hexokinase method and HbA1c was measured by high performance liquid chromatographic (HPLC) method (Bio-RAD D-10, Bio-RAD, Hercules, CA, USA).

All 30 patients had been maintained on Liraglutide in a dose of 1.2 mg per day and all 30 patients on dulaglutide 1.5 mg dose once-weekly during the thirteen weeks of the study period. Both the arms were well matched as far as baseline characteristics were concerned. (Table 2 – Supplementary appendix).

Statistical Methods:

Descriptive statistical analysis were carried out with a Statistical software named SAS (Statistical Analysis System) version 9.2 for windows, SAS Institute Inc. Cary, NC, USA and Statistical Package for Social Sciences (SPSS Complex Samples) Version 21.0 for windows, SPSS, Inc., Chicago, IL, USA, with Microsoft Word and Excel being used to generate graphs and tables. Results on continuous measurements are presented as Mean ± SEM and results on categorical measurements are presented in Number (%). Significance is assessed at a level of 5 %.

The following assumptions on data were made: 1) Cases of the samples should be independent, 2) The populations from which the samples are drawn have the same variance (or standard deviation) and 3) The samples drawn from different populations are random.

Normality of data was tested by Anderson Darling test, Shapiro-Wilk, Kolmogorov-Smirnoff test and visually by QQ plot. Paired t-test was used to find the significance of study parameters within groups of patients measured on two occasions. Chi-square/ Fisher Exact test has been used to find the significance of study parameters on categorical scale between two or more groups.

Results:

Impact of adding either Dulaglutide or Liraglutide to Metformin and SGLT2 Inhibitors:

Dulaglutide:

Adding dulaglutide to the combination of metformin 2000mg/day and aSGLT2i resulted in a significant reduction in FPG (- 41.87 ± 12.72 mg/dL; p=0.003) and HBA1C at 3 months follow up (-1.017 ± 0.22%; p<0.001). [Table 1] The impressive reduction in glycemic parameters was accompanied by a significant impact on body weight (-4.20 ± 0.47 kg; p<0.001 and BMI (-1.53± 0.21%; p<0.001) without an impact on BP.

Liraglutide:

As with dulaglutide, treatment with liraglutide in addition to Metformin and a SGLT2i resulted in a significant reduction in FPG (-44.20± 8.05 mg/dL; p<0.001), HBA1C (1.547± 0.22%; p<0.001), weight (-5.83 ± 0.87 kg; p<0.001) and BMI (2.27± 0.33&; p<0.001) at 3 months follow up. [Table 3] However in contrast to dulaglutide, liraglutide was associated with a significant reduction in SBP (-10.23 ± 2.36 mm of Hg; p<0.001). [Table 1]

References:

(1). DeFronzo R.A. From the Triumvirate to the Ominous Octet: A New Paradigm for the Treatment of Type 2 Diabetes Mellitus. Diabetes 2009 Apr; 58(4): 773-795. (2). Abdul-Ghani M.A, Puckett C, Triplitt C, Maggs D, Adams J, CersosimoE&DeFronzo R.A. Initial combination therapy with metformin, pioglitazone and exenatide more effective than sequential add-on therapy in subjects with new-onset diabetes. Results from the Efficacy and Durability of Initial Combination Therapy for Type 2 Diabetes (EDICT): a randomized trial. Diabetes, Obesity and Metabolism 2015; 17: 268–275. (3). Zinman B, Wanner C, LachinJM, Fitchett D, Bluhmki E, Hantel S, et al. Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes.NEngl J Med 2015; 373:2117-28. (4). Marso SP, Daniels GH, Brown-FrandsenK,Kristensen P, Mann JFE,Nauck M, et al. Liraglutide and Cardiovascular Outcomes in Type 2 Diabetes. N Engl J Med 2016; 375:311-322. (5). Marso SP, Bain SC, ConsoliA,Eliaaschewitz FG, JoA&Ldar E, Leiter LA, et al. Semaglutide and Cardiovascular Outcomes in Patients with Type 2 Diabetes. N Engl J Med 2016; 375:1834-1844. (6). Frias JP, Guja C, Hardy E, Ahmed A, Dong F, Öhman P, et al. Exenatide once weekly plus dapagliflozin once daily versus exenatide or dapagliflozin alone in patients with type 2diabetes inadequately controlled with metformin monotherapy (DURATION-8): a 28 week, multicentre, double-blind, phase 3, randomised controlled trial. Lancet Diabetes Endocrinol2016; 4: 1004–16. (7). Nauck AM & Meier J. GLP-1 receptor agonists and SGLT2 inhibitors: a couple at last? The Lancet Diabetes and Endocrinology 2016; 4(12): 963-964. (8). Saroka RM, Kane MP, Busch RS, Watsky J, Hamilton RA. SGLT-2 inhibitor therapy added to GLP-1 agonist therapy in the management of type 2 diabetes. EndocrPract. 2015; 21:1315-1322. (9). Curtis L, Humayun MA, Walker J, Hampton K, Partridge H. Addition of SGLT2 inhibitor to GLP-1 agonist therapy in people with type 2 diabetes andsuboptimal glycaemic control. Practical Diabetes 2016; 33(4): 129–132. (10). Association of the magnitude of weight loss and changes in physical fitness with long-term cardiovascular disease outcomes in overweight or obese people with type 2 diabetes: a post-hoc analysis of theLook AHEAD randomised clinical trial. The Look AHEAD Research Group.Lancet Diabetes Endocrinol. 2016; (11):913-921.

Dulaglutide versus Liraglutide failing OHA regime:

Both dulaglutide and liraglutide were independently associated with a significant impact on HBA1C and weight with liraglutide having an additional impact on SBP. We also analyzed the data to compare the effects of dulaglutide with liraglutide, when they were both added to existing therapy of Metformin and SGLT2i.

Glycemic Parameters:

Compared to dulaglutide, liraglutide was associated with a non-significantly greater reduction in both FPG (-44.2±8.05 mg/dL with liraglutide versus -41.87 ± 12.72 mg/dL with dulaglutide) and HBA1C (-1.55 ± 0.22 with Liraglutide versus -1.02 ± 0.22 with dulaglutide) (p= 0.091). [Table 3] However, 63.3% of patients in the liraglutide achieved an HBA1C of <7.0% compared to only 30% in the dulaglutide arm. This difference was statistically significant (0.019). [Table 3 – Supplementary appendix]

TABLE 3- Change in study parameters during the follow-up period								
Cohort	Dulaglutide, n=30				Liraglutide, n=30			
Parameter	Baseline Mean±SEM	Follow-up Mean±SEM	Change Mean ±SEM	P	Baseline Mean±SEM	Follow-up Mean±SEM	Change Mean ±SEM	P
Body weight (kg)	87.17 ± 2.16	82.97 ± 2.05	-4.20 ± 0.47	<0.001	89.43 ± 2.60	83.60 ± 2.35	-5.83 ± 0.87	<0.001
BMI (kg/m²)	32.84 ± 0.77	31.312 ± 0.68	-1.53 ± 0.21	<0.001	34.92 ± 0.868	32.650 ± 0.77	-2.27 ± 0.33	<0.001
SBP (mmHg)	134.30 ± 2.71	130.87 ± 2.49	-3.43 ± 2.97	0.258	137.83 ± 2.503	127.60 ± 2.07	-10.23 ± 2.36	<0.001
DBP (mmHg)	80.73 ± 1.64	78.43 ± 0.914	-2.30 ± 1.79	0.208	83.47 ± 1.587	80.67 ± 1.51	-2.80 ± 1.85	0.141
FPG(mg/dl)	175.53 ± 11.45	133.67 ± 6.90	-41.87 ± 12.72	0.003	160.13 ± 8.17	115.93 ± 5.69	-44.20 ± 8.05	<0.001
HbA1c (%)	8.43 ± 0.21	7.411 ± 0.15	-1.017 ± 0.22	<0.001	8.49 ± 0.26	6.95 ± 0.21	-1.547 ± 0.22	<0.001

Weight:

There was a greater weight loss (non significant) with liraglutide (-5.83 ± 0.87 kg with liraglutide vs. -4.2 ± 0.46 kg with dulaglutide) when compared to dulaglutide (p=0.103). [Table 4 – Supplementary appendix] A similar proportion of patients in both the arms experienced a weight loss between 5-10% (53.3% with dulaglutide vs. 50.0% with liraglutide). [Table 5 – Supplementary appendix] A larger proportion of patients in the liraglutide arm (13.4%) achieved a weight loss more than 10% from baseline compared to dulaglutide (3.3%). A highly significant weight loss of greater than15% from baseline was seen in 6.7% patients in the liraglutide arm, but not in the dulaglutide arm. [Figure 1 – Supplementary appendix]

Blood Pressure:

There was similar reduction in DBP in both the arms (-2.8 ±1.85 mm of Hg with liraglutide vs. -2.3 ± 1.79 mm of Hg with dulaglutide (p=0.847). [Table 4 –Supplementary appendix] In contrast the reduction in SBP was greater with liraglutide (-10.23 ± 2.36 mm of Hg) compared to dulaglutide (-3.43 ± 2.98 mm of Hg) (p=0.079), but failed to reach statistical significance.

Dulaglutide vs. Liraglutide: Impact on the metabolic composite of HBA1C reduction AND weight loss:

The composite end point of HBA1C reduction to below 7% and a greater than 5% of body weight loss was attained by 5 (16.7%) patients in the dulaglutide arm, while 20 (50%) patients on liraglutide arm reached this composite. This difference was not statistically significant. (p = 0.75). (Figure 1)

Adverse Events:

No serious adverse event was reported during this three month follow up. There was no hypoglycaemia reported. The commonest adverse event was nausea in 14 (23%) patients, 6 (20%) in the dulaglutide group compared to 8 (26.6%) in the liraglutide group. Two patients complained of genital irritation, which responded to topical anti mycotics.

Discussion:

The mode of therapy co-initiating SGLT2i and GLP1RA was recently studied and published as the DURATION 8 study showing a robust weight loss and reduction in SBP in the dapagliflozin plus Exenatide-LAR arm. Strangely, this benefit did not extend to HBA1c reduction. (6) It was speculated that the rapid glucose lowering effect of SGLT-2 inhibitor could have blunted the robust glucose lowering effect of GLP-1 RA due to the co initiation of the 2 molecules. (7) Hence it would be interesting to take a look at the effect on glycaemic parameters when these two agents are added sequentially.

A couple of real-world observational data did find additive benefit of combining SGLT-2i to GLP1-RA. (8,9)

In this real world study from India, we looked at the more realistic scenario of GLP1RA therapy being instituted when oral medicines have been suboptimal and this is the first data using this combination in this order.

Both liraglutide as well as dulaglutide resulted in a significant reduction in FPG, HBA1C as well as weight from baseline. Only liraglutide not dulaglutide had a significant impact on the reduction of SBP. It should be noted here that all patients in this cohort had a reasonably well-controlled blood pressure at baseline and no antihypertensive medications were changed during the study.

This data would therefore suggest that as a first injectable, GLP1-RA is a very effective option when it is used along with a SGLT-2 inhibitor on background metformin therapy in a sequential approach.

Our study also looked at the comparative effectiveness of liraglutide versus dulaglutide as an add on option to a SGLT-2 inhibitor and metformin. There seems to be a trend favoring liraglutide vs. dulaglutide in reduction of fasting plasma glucose, HBA1C, weight and SBP. There was a statistically significant difference favoring liraglutide (63.3% with liraglutide and 30% with dulaglutide) in the proportion of patients achieving HBA1C less than 7.0%. Larger proportion of patients achieved target HBA1C without weight gain while on liraglutide.

Off note, no serious adverse events were noted during the study.

There are several limitations in this study. Firstly it was open label and not controlled. Hence this data could not correct for numerous confounding factors that could have influenced the results. Secondly the small sample size could have influenced the quantum of metabolic impact. We are aware that this size was small also because only a small percentage of our patients could afford this very expensive therapy in our “pay from pocket” setting and were “chosen” based on their ability to afford this treatment. This would also be considered a major selection bias.

Conclusion:

In this retrospective study in Indian type 2 diabetic patients poorly controlled with metformin and SGLT-2 inhibitor we found a meaningful impact of adding a GLP1-RA on all metabolic parameters. Both liraglutide and dulaglutide were effective in this regard. There were additional advantages seen with liraglutide as far achieving target HBA1C of less than 7% and also on the quantum of weight loss and SBP reduction.

This study therefore provides a pilot and indeed generates a hypothesis supporting the sequential addition of a GLP1-RA after metformin and SGLT-2 inhibitor, in the management of T2D, which needs to be tested in a more systematic manner on a larger population.