HEALTH-RELATED QUALITY OF LIFE OF BREAST CANCER PATIENTS, FIVE-YEAR FOLLOW-UP

Roine E¹, Saarto T¹, Blomqvist C¹, Sintonen H²

1. Helsinki University Hospital, Comprehensive Cancer Center, Helsinki, Finland
2. Department of Public Health, University of Helsinki, Helsinki, Finland

INTRODUCTION

- Adjuvant treatments of breast cancer (BC) improve survival but are associated with adverse effects.
- Measuring health-related quality of life (HRQoL) gives information on the patients' own views of their subjective well-being.

OBJECTIVES

- To measure HRQoL of BC patients and to compare their HRQoL to that of the general population.

METHODS

- 573 BC patients aged 35-68 (mean 53) recruited to an exercise intervention study were after adjuvant treatment randomly allocated into an intervention or control group.
- The patients filled in the 15D generic HRQoL questionnaire at baseline and at 5 years follow-up.
- The HRQoL of the patients was compared to that of the age-standardized general female population.

RESULTS

- Compared to the general population, the BC patients (n=181) had statistically significantly and clinically importantly worse mean HRQoL score at baseline (Fig.1).
- At five-year follow-up, the BC patients' (n=390) mean HRQoL score had improved but was still statistically significantly and clinically importantly worse than that of the general population (Fig.2).
- The most significant differences (p<0.01) were on the dimensions of sleep, excretion, mental function, distress, vitality and sexual activity (Fig.2).
- The impairment in HRQoL score was more pronounced in younger patients (<53 years at baseline, n=191) at both time points (Fig.3).
- In the general population HRQoL scores tend to decrease with rising age. In the BC patients the opposite was seen with older BC patients (>52 years at baseline, n=216) having better mean HRQoL scores than younger patients at both time points.

CONCLUSIONS

- Five years after the start of treatment, BC patients still had worse mean HRQoL score than the general population.
- The difference was greatest in younger patients.

Figure 1. Comparison to population at baseline.

Figure 2. Comparison to population at five-year follow-up.

Figure 3. Comparison to population at five-year follow-up in younger (<53 years) patients.