ANALYSIS OF NEOADJUVANT THERAPY IN VULVAR MALIGNANT NEOPLASM

L. Elias¹, M. Komatsu,¹ C. Martin Gomez¹, C. de Moraes Gomes Lopes¹, T. Gomes de Almeida¹.

¹Hospital Santa Marcelina, Ginecologia Oncológica, São Paulo, Brazil.

OBJECTIVES

Evaluate the clinical results of patients with locally advanced vulvar malignant neoplasm treated with neoadjuvant radiotherapy or chemotherapy.

METHODS

Retrospective analysis of the patients medical records who were followed up at the Gynecology Oncology department of Santa Marcelina Hospital in São Paulo between 2008 and 2018 that underwent neoadjuvant radiotherapy or chemotherapy for vulvar neoplasm.

RESULTS

In the studied period, 55 patients were diagnosed with vulvar neoplasm, 21 (38.18%) submitted to neoadjuvant radiotherapy and 15 (27.27%) underwent neoadjuvant chemotherapy too. Of the 21 patients treated with neoadjuvant therapy, 1 had histopathological diagnosis of adenocarcinoma and the other 20 of squamous cell carcinoma. Twelve patients (57.14%) underwent surgery afterwards: 10 patients (47.61%) had a radical vulvectomy with bilateral lymphadenectomy and 2 patients (9.52%) had a hemivulvectomy with bilateral lymphadenectomy. In the follow-up, 4 patients (19%) presented persistence of disease and 5 (23.8%) local recurrence. Of the patients submitted to neoadjuvant therapy, 11 (52.38%) died and 7 (33.3%) lost follow-up. The majority of patients were in stage II (FIGO 2009).

CONCLUSIONS

The evolution of vulvar neoplasm results in the involvement of structures close to the vulva, like urethra and anal region.

Thus, as most patients at the time of diagnosis already have a locally advanced disease, neoadjuvant therapy decreases the tumor load and reduces the need for extensive surgeries, also decreasing surgical morbidity. In this study, the complete control rate was approximately 57.2% with neoadjuvant therapy, demonstrating the benefit of this type of treatment.

Figure 1. Vulvar carcinoma before neoadjuvant radiotherapy and chemotherapy.

Figure 2. Macroscopic aspect of neoplasm after neoadjuvant radiotherapy and chemotherapy.

Figure 3. Macroscopic aspect of vulvar carcinoma after neoadjuvant therapy.

REFERENCES

1. Coelho F.R.G., Costa R.L.R., Almeida T.G., Mesquita J. Revisão em Ginecologia Oncológica. Ed. Andrei. 2017. 1ª Ed.
2. D.M. Forner, P. Mallmann. Neoadjuvant and definitive chemotherapy or chemoradiation for stage III and IV vulvar cancer: a pooled reanalysis. Eur J Obstet Gynecol Reprod Biol, 212 (2017), pp. 115-118.
3. Beriwal S, Shukla G, Shinde A, et al. Preoperative intensity modulated radiation therapy and chemotherapy for locally advanced vulvar carcinoma: analysis of pattern of relapse. Int J Radiat Oncol Biol Phys. 2013; 85: 1269- 1274.
4. Niu Y, Yin R, Wang D, Li Q, Gao X, Huang M. Clinical analysis of neoadjuvant chemotherapy in patients with advanced vulvar cancer: A STROBE-compliant article. *Medicine (Baltimore)*. 2018;97(34):e11786.