HARRIS HIP SCORE AND REHABILITATION IN SURGICALLY TREATED PERTROCHANTERIC FRACTURE

Mitrevska B¹, Nikolik- Dimitrova E¹, Koevska V¹, Gerakaroska-Savevska C¹,Gocevska M¹, Kalcovska- Ivanovska B¹, Manoleva M¹,Grkova-Miskovska E ²

¹Institute of Physical Medicine and Rehabilitation – Skopje, Macedonia
² Sity Hospital 8-th September, Skopje, Macedonia

Intruduction:By increasing average length of life, solving fractures of the upper end of the femur is a major problem for the modern surgery.

Purpose: To evaluate the effects of treatment and physical rehabilitation with kinesiherapy and magnetictherapy versus kinesiotherapy and therapy with interference currents of patients with surgically treated pertrochanteric fracture with DHS, according to the protocol results monitoring.

Method: The study represents a prospective randomized clinical trial . Include two cohorts, with 90 participants with surgically treated pertrochanteric fracture with DHS. Respondents are divided into two groups: Examined cohort - 45 patients is treated with kinesitherapy and magnetictherapy and control group - which has 45 patients treated with kinesitherapy and therapy with interference currents. Respondents were followed for one year, during which were performed three examinations, from the first review which is input for selected patients who meet the criteria for inclusion in research.

Results: It is recognized that, kinesitherapy and magnetictherapy for p <0,05, increase Harris hip score (improve the condition of patients) in three time combinations. For p <0,05, significant difference with Wilcoxon signed rank test was perceived between Harris hip score at: a) outs / 6 months Z = 5,712 p = 0,0001 in addition to 6 months; b) discharge / 12 months Z = 5,842 p = 0,0001 in addition to 12 months; c) 6 months / 12 months Z = 5,844 p = 0,0001 in addition to 12 months. It is recognized that, kinesitherapy and interference currents, for p <0,05, increase Harris hip score (improve the condition of patients) in three time combinations. For p <0,05, significant difference with Wilcoxon signed rank test was perceived between Harris hip score : a) discharge / 6 months Z = 5,842 p = 0,0001 in addition to 6 months; b) discharge / 12 months Z = 5,842 p = 0,0001 in addition to 12 months; c) 6 months / 12 months Z = 5,778 p = 0,0001 in addition to 12 months.

Comparison of the two groups after Harris hip score

times of physical examination

	discharge	6 mounths	12 mounths
Mann- Whitney U	620	451	385
Z	(3.168)	(4.535)	(5.085)
Asymp.Sig. (2-tailed)	0.002*	0.0001*	0.0001*

Upon discharge, after 6 and 12 months, for p< 0,05, the analysis indicated a statistically significant difference between the two groups in terms of Harris hip score

Analysis of the examinated group of Harris hip score for three times

For p <0,05, in the examined group found a significant difference between Harris hip score in the three times of physical examination. It is recognized that, kinesitherapy and magnetictherapy for p <0,05, increase Harris hip score (improve the condition of patients) in three time combinations.

Conclusions: On physical examination after discharge, after 6 and 12 months, Harris hip score in the examined group with magnetictherapy was significantly greater and the condition of patients was significantly better compared, with the same, in the group treated with kinesitherapy and interference currents.

It can be concluded that postoperative rehabilitation of pertrochanteric fractures with fixation of dynamic implant-DHS, therapy of choice is kinesiotherapy and magnetictherapy, from which there is an improvement both in functional status, in the stimulation of osteogenesis and quality of life in adult patients.

According to the comparison of median Harris hip score in both groups, analysis indicated that the situation after 12 months in 50% of patients treated with kinesitherapy and magnetictherapy are assessed as **excellent**, compared to the 50% of patients treated with kinesitherapy and interference currents which are assessed as rather **Good**