Current Practice of Preoperative Fasting for Pediatric Day Case Surgery in a tertiary care university hospital in Egypt

Nassar H., Fawzy M., El-Hadi H., Al-Hakami M

Cairo University, Dept of Anaesthesiology & Intensive Care, Cairo, Egypt

Background and Goal of Study:

Pediatric fasting guidelines are designed to reduce patient discomfort while minimizing the risk of perioperative morbidities. This study was designed to determine current practice of pediatric preoperative fasting for day case surgeries in our Hospital, effect of prolonged preoperative fasting on perioperative morbidities and to assess the possible role of doctor/parent / hospital interactions in changing the current status.

Materials and methods:

This cross-sectional study was conducted in Cairo University paediatric hospital from June to October 2016. Participants were children (2 months -12 years, ASA I or II)undergoing elective day case surgery under general anesthesia; their parents and pediatric surgery residents (SR) participated in questionnaires. We focused on the actual preoperative fasting duration(primary outcome), perioperative morbidities, parents understanding and compliance with fasting instructions as well as on the SR familiarity and acceptance of fasting guideline recommendations

Results and discussion:

200 children (mean age 4 years; ophthalmological surgeries 57.5%) and their parents (200 mothers) were enrolled in this study as well as 16 SR. Our children experienced prolonged fasting before procedures compared to ASA recommendation times(p 0.001) with no incidence of regurgitation or pulmonary aspiration(mean fasting times for oral fluids, breast milk, formula and solids were 7.28(1.49), 5.64(1.13),6.86(0.97),7.95(2.7) hours respectively). Blood sugar level was less than 70 mg/dl in 8 cases, 75.0% of them were fasting for more than 8 hours (figure 1). Only18.5% of the parents believed that fasting was necessary to avoid the development of health problems. All parents (100%) and 80% of SR didn't know that preoperative fasting ASA guidelines allow oral clear fluids up to 2 hours preoperatively (figure 2). Concerns regarding patient safety and every day work flow in our institution might explain the reasons against implementing fasting guidelines.

Figure 1:Random blood sugar categories at different fasting duration and age groups

Figure 2:preoperative fasting hours recommended by residents

Conclusion(s):

Prolonged pediatric preoperative fasting period for day case surgery is still a common practice in our institution. Hypoglycemia incidence increases with fasting more than 8 hours. Knowledge deficit about fasting guidelines among our parents and SR as well as concerns regarding patient safety and every day work flow in our institution were the major impediments for improving preoperative fasting care .