Gastrectomy with extended lymphadenectomy: a North-American perspective

Alexandre Gosselin-Tardif¹, Jessica Lie¹, Ioana Nicolau², Jonathan Cools-Lartigue², Liane Feldman¹, Jonathan Spicer², Carmen Mueller^{1,2}, and Lorenzo Ferri²

Department of General Surgery¹, Division of Thoracic Surgery², McGill University Health Centre, Montreal, QC Canada

Introduction

- Ongoing debate on extent of lymphadenectomy in gastric adenoCa
- Increasing evidence of long-term survival benefit with extended (D2) lymphadenectomy
- Despite this, concerns over post-op complications from D2 lead many N-A surgeons to perform traditional (D1) node dissection

Purpose

- To evaluate the **safety and efficacy of D2 dissection** as performed in a high-volume North American center

Methods

- Retrospective, single-institution cohort study at McGill University Health Centre (MUHC) of all non-bariatric gastrectomies, 2005-16.
- Patients grouped D1 vs D2, comparing operative outcomes
- Subgroup analysis of laparoscopic vs open D2
- **1° outcomes:** Postoperative morbidity and mortality, anastomotic leak rates, nodal yield, resection margin status
- Exclusion criteria: Wedge resection, thoracoabdominal approach, emergency operation, palliative surgery

FIGURE 1 – Patient selection

TABLE 1 – Patient and treatment characteristics

	D1 n=55	D2 n=151	p-value
Age in years	71.8 (58-80)	70.1 (60-77)	0.76
Female gender	20 (36)	51 (34)	0.73
2+ comorbidities	25 (46)	62 (41)	0.69
Body mass index	23.2 (4.6)	25.2 (5.3)	0.20
Indication			<0.001
- Adenocarcinoma	34 (62)	141 (93)	
- Other tumor	7 (13)	6 (4)	
- Benign disease	10 (18)	1 (1)	
- Prophylactic	4 (7)	3 (2)	
Procedure			0.020
- Total	13 (24)	56 (37)	
- Distal	40 (73)	78 (52)	
- Proximal	2 (4)	17 (11)	
Laparoscopic	23 (42)	48 (32)	0.27
Splen/panc-ctomy	1 (2)	9 (6)	0.29
Neoadjuvant CTX	2 (4)	55 (36)	<0.001
Adjuvant CTX	9 (16)	67 (44)	<0.001
Stage (n=175)			<0.001
- Stage 1	21 (62)	40 (29)	
- Stage 2	2 (6)	41 (29)	
- Stage 3	11 (32)	60 (43)	

Among D2, higher proportion of ADK as indication, total gastrectomy, high stage cancer and CTX.

All data presented as n(%) or median (IQR)

Results

TABLE 2 – Operative outcomes

	D1 n=55	D2 n=151	p-value
Operative length, min	197 (98)	210 (70)	0.25
Lymph node yield	13 (7-22)	30 (23-42)	<0.001
R0 resection (n=175)	33 (97.9)	129 (91.5)	0.47
All complications	22 (40)	61 (40)	0.96
Severe complications (grade >2)	12 (22)	25 (17)	0.38
Mortality	3 (5)	3 (2)	0.19
Anastomotic leak	4 (7)	8 (5)	0.74
Intra-abdominal abscess	2 (4)	8 (5)	1.0
Re-operation	7 (13)	15 (10)	0.57
Length of stay, days	8 (6)	8 (7)	0.18
30-day readmission	6 (11)	13 (9)	0.61

No difference in morbidity, mortality, leaks or R0 resection. Higher nodal yield after D2 resection.

TABLE 3 – Laparoscopic vs Open D2 dissection

	Lap D2 n=48	Open D2 n=103	p-value
Body mass index	25.7 (4.7)	24.0 (5)	0.085
Stage (n=141)			0.015
- Stage 1	20 (44)	20 (21)	
- Stage 2	10 (22)	31 (32)	
- Stage 3	15 (33)	45 (47)	
Operative length, minutes	199 (61)	212 (65)	0.15
Lymph node yield	31 (13)	29 (23)	0.79
R0 resection (n=141)	41 (91)	88 (92)	1.0
All complications	21 (44)	40 (39)	0.57
Anastomotic leak	2 (4.2)	6 (5.8)	1.0
Re-operation	7 (14)	8 (8)	0.19
Length of stay, days	5 (6)	9 (7)	<0.001

No difference in complications, nodal yield, R0 resection between lap and open D2. Shorter LOS, lower stage associated with lap D2.

Discussion

- Study highlights safety and efficacy of D2 in North American setting, with superior lymph node yield to D1 achieved at equivalent complication rates.
- Also highlights feasibility of lap D2 in North American body habitus, with equivalent perioperative and oncologic results, as well as reduced LOS.

Limitations

- Retrospective design
- Type of lymphadenectomy influenced by surgeon preference, patient and tumor characteristics.
- Outcomes may be influenced by differences in surgical indication, type of gastrectomy.

We would like to thank the members of the Division of Thoracic Surgery of the Montreal General Hospital for their support, as well as the Canadian Association of General Surgeons for the opportunity to present our findings