“Prevalence of MYD88 L265P mutation in Diffuse Large B-Cell Lymphomas involving the skin: Single-center study in Argentina.”

Dana Kohan, Federico Jauk, Julieta Pandolfi, Victoria Volonteri, Victoria Otero, Hernán García Rivello.
Pathology Department, Italian Hospital of Buenos Aires, Argentine

Introduction: Cutaneous involvement by diffuse large B-cell lymphoma (DLBCL) can occur primarily in the skin, or secondarily to a systemic DLBCL (S-DLBCL). In the first group, two different clinicopathological entities can be separated: primary cutaneous large B-cell lymphoma not otherwise specified (PCLBCL-NOS), and primary cutaneous large B-cell lymphoma leg-type (PCLBCL-LT). MYD88 L265P mutation was proven to be present in 16,5% of systemic DLBCL and 40 - 68% of PCLBCL-LT cases, and is associated with poor prognosis and shorter overall survival.
Aim: Determine the prevalence of MYD88 L265P mutation in PCLBCL-NOS, PCLBCL-LT and S-DLBCL.

Materials and Methods

A retrospective observational study was performed. All cutaneous DLBCL cases diagnosed in Italian Hospital of Buenos Aires (Argentina) between 2007 and 2017 were included. 15 cases were selected and reclassified based on the WHO criteria (2017) in three groups: S-DLBCL (n=6), PCLBCL-NOS (n=6) and PCLBCL-LT (n=3).

If necessary to reclassify cases following the WHO (2017) criteria, immunohistochemistry (IHC) was performed, using an automated processor (Benchmark XT, Ventana). DNA was extracted from formalin-fixed paraffin-embedded tissue, using QIAamp DNA mini kit (QIAGEN). The presence of MYD88 L265P was assessed using qBiomarker Somatic Mutation PCR Assay Kit, specific for c.794T>C substitution in MYD88 gene.

Results:

Table 1.

	Age (median; range)	Sex (Male/Female)
PCLBCL-LT	84; 79-85	2/1
PCLBCL-NOS	69; 65-85	4/2
S-DLBCL	73; 59-80	6/0

Table 1. Clinical features of the three groups of patients.

Table 2. Location, immunophenotypic profile and MYD88 status in all patients involved in this study.

Table 2.

	Case no.	Location	CD20	CD10	Bcl6	MUM1	Bcl2	MYD88
PCLBCL-LT	1	Arm	Positive	Positive	Positive	Positive	Positive	Positive
	2	Leg	Positive	Positive	Positive	Positive	Positive	Positive
	3	Pectoral. Chin	Positive	Positive	Positive	Positive	Positive	Positive
PCLBCL-NOS	4	Scalp	Positive	Positive	Positive	Positive	Positive	Positive
	5	Scalp	Positive	Positive	Positive	Positive	Positive	Positive
	6	Leg	Positive	Positive	Positive	Positive	Positive	Positive
	7	Scalp	Positive	Positive	Positive	Positive	Positive	Positive
S-DLBCL	8	Zigomatic region	Positive	Positive	Positive	Positive	Positive	Positive
	9	Thigh and back	Positive	Positive	Positive	Positive	Positive	Positive
	10	Forearm	Positive	Positive	Positive	Positive	Positive	Positive
	11	Flanck	Positive	Positive	Positive	Positive	Positive	Positive
	12	Not specified	Positive	Positive	Positive	Positive	Positive	Positive
	13	Thigh	Positive	Positive	Positive	Positive	Positive	Positive
	14	Leg	Positive	Positive	Positive	Positive	Positive	Positive
	15	Neck	Positive	Positive	Positive	Positive	Positive	Positive
	16	Scalp	Positive	Positive	Positive	Positive	Positive	Positive

Figure 1. Representative case (case No X) showing morphological characteristics (A and B) and immunohistochemical profile of group PCLBCL-LT (C, D and E).

Figure 2. Amplification plot showing MYD88 reference (green line) and MYD88 L265P (red line) fluorescent signal. Case 2 (plot A) shows both reference and mutant amplification. Case 6 (plot B) shows reference amplification but not mutant amplification.

Conclusions

In our series, L265P MYD88 mutation was observed in 66,7% of PCLBCL-LT and 16,5% of S-DLBCL, in coincidence with previous reports. Although a small amount of cases were included, we plan to expand this study in a collaborative way with other institutions from our country. Testing MYD88 mutation could be useful as prognostic and predictive factor both in primary and secondary involvement of the skin by a DLBCL, but further investigation is needed.