LOW SPECIFICITY OF NICE GUIDELINE IN RECOGNITION OF PAEDIATRIC SEPSIS

S. Omar¹, L. Archer¹, R. Mills¹, F. Khurshid¹, S. Welch¹.

¹Department of Paediatrics, Heartlands Hospital, University Hospitals Birmingham NHS Foundation Trust, United Kingdom

University Hospitals
Birmingham
NHS Foundation Trust

Background

The recognition of sepsis is a significant challenge to paediatricians, with devastating consequences if missed.

In 2016 the NICE Sepsis Guidelines (NG51) were published, stratifying sepsis risk based on age, observations and examination into low, moderate and high risk.

The guidelines provide subsequent management recommendations based on the risk category.

Aim To compare actual recognition and management of suspected sepsis in children to the NICE suggested recognition and management.

Method

A retrospective audit was carried out at a busy paediatric unit in England over two weekends (February and July) in 2017. Data was collected using notes and electronic systems.

Information collected included:

- Admission observations – Heart rate, respiratory rate, blood pressure, temperature and capillary refill time
- Clinical examination
- Investigations undertaken
- Suspected and final diagnosis

Using the data and NICE guidelines the sepsis risk was stratified for each patient as low, moderate or high. The suggested NICE management pathway for that sepsis risk was then compared to the actual management of the child based on the clinician's assessment.

Results

Of 93 attendances; 64 (69%) had an admission diagnosis of infection. As per NICE, 38 (60%) of these would have stratified as high risk warranting sepsis management (including bloods, chest x-ray, antibiotics, oxygen and IV fluids).

Based on clinician assessment, only 4 were treated as sepsis. Of these the final diagnoses were:

- Sepsis secondary to UTI x 2
- Encephalitis
- Scarlet fever

Of the 34 high risk patients not treated for sepsis, 24 (63%) were diagnosed with viral illness and none developed sepsis.

Another 9 patients diagnosed with an infection had 2 moderate sepsis risk factors as per NICE but none of these were treated for or developed sepsis.

Conclusion

With regards to recognising sepsis in children the NICE guideline had a; sensitivity of 100%, specificity of 27%, positive predictive value (PPV) of 4% and negative predictive value (NPV) of 100%.

In comparison, clinician assessment had a sensitivity of 100%, specificity of 97%, PPV of 50% and NPV of 100% for recognising sepsis (figure 1).

If the guideline had been followed a further 43 children would have undergone inappropriate sepsis management (figure 2).

Take Home Message

This study demonstrates that the NICE sepsis guidelines are poorly specific for sepsis and would lead to unnecessary investigations, antibiotics and admission for a large number of patients.

Figure 1: Comparison of NICE guidelines and clinical assessment for recognition of sepsis in children

Tool used	Sensitivity	Specificity	PPV	NPV
NICE guidelines	100%	27%	4%	100%
Clinician Assessment	100%	97%	50%	100%

Figure 2: Comparison of Management in Children with Suspected Infection