THE ROLE OF VITAMIN D IN WOMEN WITH POSTMENOPAUSAL OSTEOPOROSIS

Koevska V^{1.}, Nikolik-Dimitrova E^{1.}, Mitrevska.B^{1.}, Manoleva M^{1.}, Gjerakaroska-Savevska C^{1.}, Kalcovska-Ivanovska B^{1.}, Gocevska M^{1.}

¹Institute for Physical Medicine and Rehabilitation, Skopje, R. Macedonia

Introduction: In the treatment of postmenopausal osteoporosis (PMO), vitamin D and calcium has very important role to maintain and improve bone mineral density (BMD). Women with low vitamin D values have reduces absorption of calcium from the gastrointestinal tract.

Material and methods: 92 women with PMO have regular physical activity, and takes every day 800IU vitamin D and Calcium 1000mg. The level of serum vitamin D was measured by electrochemiluminiscence,at reception and after 12 months research. BMD was determined at, the lumbar spine, L2-L4 and femoral neck by dual-energy x-ray absorptiometry, at baseline and after 12 months research.

Results: The average age of patients was 60.64 ± 6.7, education: primary 22 (23.9%), intermediate 45 (48.91%), and higher 25 (27.17%). The results have shownen statistically significant value in mean vitamin D level (tab.1) also, statistically significant value in mean BMD of the lumbar spine and femoral neck, at start and after 12 months research (lumbar spine 0.71 ± 0.2g/sm2 v.s 0.77 ± 0.1g/sm2, p<0.0001 and neck of femur 0.8 ± 0.12 g/sm2 v.s 0.83 ± 0.1 g/sm2, p = 0.001). Chart 2. and Ch.3.

Discussion: Studies have shown that there is a high percentage of low vitamin D levels in PMOS (1,2). Our research showed a high percentage of vitamin D deficiency in the blood (83%). Regular monitoring and regular vitamin D intake an increase the percentage of patients (56.52%) with normal vitamin D values after 12 months . The association of vitamin D, calcium, exercise and BMD is questionable (3).

Nevertheless, the results of the several research suggest that regular intake of vitamin D, calcium and exercises have a significant role to play in maintaining and improved BMD in PMOP (2,4).

Conclusion: Our sample of research, although modest, has shown that vitamin D, calcium and exercise plays a significant role in maintaining and enhancing BMD in PMOP.

Table 1. Statistical difference of the average total vitamin D at admission and after 12 months

	Descriptive statistics (vitamin D/ µg)		p value
	mean±SD	min – max	
Reception	20,4 ± 9.03	6,59 – 47.0	t=10.11
Control	30.93 ± 7.95	8.85 – 45.0	p<0.0001

Table 2. Distribution of patients according to the level of vitamin D in the blood; deficiency, insufficiency and normal vitamin D.

vitamin D / ng	n (%)		
	deficiency	insufficiency	normal
Reception	13 (14.13)	70 (76.09)	9 (9.78)
Control	2 (2.17)	38 (41.3)	52 (56.52)

Chi-square = 47.9
p<0.001

Chart 2.

Chart 3.

1. Gaugris RP et al. Vitamin D inadequacy among post-menopausal women: a systematic review. Q J Med. 2005; 98:667–676.
2. Labronici PJ et al. Vitamin D and its Relation to Bone Mineral Density in Postmenopause Women. Rev. Bras. Ortop. 2013;48(3):228-35
3. Kamineni, et al.: Serum vitamin D levels and bone mineral density in normal postmenopausal women Journal of Mid-life Health, 2016;7:4.
4. Kemmler W, Engelke K, and Stengel SVLong-Term Exercise and Bone Mineral Density Changesin Postmenopausal Women—Are There Periods of Reduced Effectiveness? Journal of Bone and Mineral Research, 2016, 31(1)p. 215–222.