Risk factors involved in the onset of postoperative delirium after transurethral prostatectomy in elderly- a retrospective study

Savu Carmen¹, Burchiu Eliza¹, Surcel C^{2,3}, Mirvald C^{2,3}, Mihai V², Pavelescu C², Tulbure Ruxanda¹

¹Anesthesiology and Intensive Care Unit, Fundeni Clinical Institute, Bucharest, Romania

²Centre of Uronephrology and Renal Transplantation, Fundeni Clinical Institute, Bucharest, Romania.

³“Carol Davila” University of Medicine and Pharmacy, Bucharest, Romania.

Introduction:

- The **aim** of this study was to investigate the incidence of postoperative delirium (POD) in elderly patients undergoing monopolar TURP for benign prostatic hyperplasia (BPH) and to identify the risk factors involved in the onset and severity of POD.
- The elderly represent a large cohort of patients undergoing urological procedures in which transurethral prostate resection (TURP) is one of the most frequently performed.

Material and method:

- 240 patients aged over 70 years underwent elective surgery for BPH in our Department between Jan 2013- Dec 2015.
- In all cases, monopolar transurethral prostatectomy (TURP) was performed. Distilled water was used as rinsing liquid.
- Patients were evaluated preoperatively, immediately postoperative and at 5 days after TURP.
- Inclusion criteria:
 - spinal anesthesia;
 - preoperative minimal state examination (MMSE)>27.
- Patients with psychiatric disorders, preoperative antidepressants, or who developed postTUR syndrome were excluded.
- Postoperative delirium was diagnosed using the Confusion Assessment method (CAM).
- After surgery, patients were divided into 3 groups based on POD severity:
 - Group 1- no delirium – 217 (90.41%) patients
 - Group 2 – POD for 2-3 days – 17 (7.08%) patients
 - Group 3 – POD >4 days – 10 (4.16%) patients
- Perioperative data included operating time, TAM, HR, SpO2, rectal temperature, blood loss, volume of rinsing liquid.
- POD treatment consisted in titrated Haloperidol (0.5 mg i.v at a time); pain management and non-pharmacological measures: early nutrition, assist with the sleep-wake cycle, presence of family at the bedside, glasses and hearing aid, etc.
- Benzodiazepines were used if sedation was required.
- For comparison between these 3 groups we used Mann-Whitney U test for continuous data, Kruskal Wallis test for non parametric variables and χ^2 test for ordinal data. A p value<0.05 was considered statistically significant.

Results:

Demographic data	Total	Group 1	Group 2	Group 3	p
Age (mean, CI)	76.3 (70-87)	75.2 (70-87)	76.1 (71-81)	74.8 (70-85)	0.474
Prostate volume, cc (mean, CI)	65.22 (25.6-85.4)	60.21 (26.33-76.51)	62.47 (25.6-81.44)	64.21 (26.6-85.4)	0.321
History of ↑ alcohol intake No. (%)	54 (22.5)	32 (15.02)	11 (64.7)	6 (60)	<0.001
History of smoking No. (%)	73 (30.41)	24 (11.26)	6 (35.29)	4 (40)	0.032
Preop. urinary catheter No. (%)	14 (5.83)	3 (1.4)	7 (41.17)	4 (40)	<0.001
ASA score (mean, CI)	3 (2-4)	3 (2-4)	3 (2-4)	3 (2-4)	0.214
Marital status - single No (%)	31 (12.91)	12 (5.63)	12 (70.58)	7 (70)	0.032
Preoperative Hb (mean, CI)	11.4 (8.75-15.17)	11.1 (9.44-14.78)	10.78 (8.75-13.45)	11.24 (9.31-15.17)	0.421
Previous surgery (No, %)	28 (11.6)	19 (8.92)	6 (35.29)	2 (20)	0.15
Preop MMSE (points), mean	28 (27-30)	29 (27-30)	28 (27-30)	28 (27-30)	0.345

Perioperative data	Total	Group 1	Group 2	Group 3	p
Operative time, min, (mean, CI)	56.21 (35-104)	48.35 (42-71)	61.14 (35-94)	58.22 (41-104)	0.042
Mean Rectal temperature (° C)	36.8 (35.1-37.2)	37 (35.1-37.2)	36.8 (35.1-37.2)	36.6 (35.1-37.2)	0.032
Hypothermia time, min, (mean, CI)	39.64 (20-84)	33.14 (20-41)	36.45 (20-74)	52.36 (30-84)	0.005
Rinsing liquid (L), mean , CI	18.21 (12.2-38.6)	17.54 (12.4-27.2)	19.75 (13.1-37.8)	18.08 (12.2-38.6)	0.122
Blood loss (g), mean, CI	2.14 (0.4-4.21)	1.61 (0.4-1.92)	2.41 (1.1-3.6)	2.12 (0.6-4.21)	0.032

Postoperative data	Total	Group 1	Group 2	Group 3	p
Hospital stay (days) mean, CI	7.1 (3-21)	5.4 (3-10)	7.3 (5-12)	12.6 (6-21)	<0.004
ICU stay (days) mean, CI	3 (0-7)	1 (0-1)	3.14 (2-4)	6.51 (4-7)	<0.004
Complication rates Clavien scale (No,%)					
I	29 (12.08)	20 (9.38)	7 (41.17)	2 (20)	0.24
II	27(11.25)	0 (0)	17 (100)	10 (100)	0.1
IIla	5 (2.08)	0 (0)	4 (23.52)	1 (10)	0.98
>IIla	0 (0)	0 (0)	0 (0)	0 (0)	n/a
Blood Transfusion No (%)	2 (0.83)	0 (0)	1 (5.82)	1 (10)	n/a

	Univariate			Multivariate		
	HR	95% CI	p	HR	95% CI	p
Age (continuous variable)						
<76 (reference)	1.000			1.000		
>76	2.049	1.722–2.596	<0.0001	1.794	1.7255–2.8999	<0.0001
Marital status-single						
No	1.266	1.187–1.351	0.04	1.2331	0.610–1.302	0.018
Yes	3.247	2.703–3.900	<0.0001	2.5963	1.9142–3.5214	<0.0001
History of ↑ alcohol intake	1.23	1.159–1.575	0.004	1.6279	1.207–2.196	0.003
Preoperative urinary catheterization	1.914	1.637–2.438	<0.0001	1.4172	1.1946–1.6814	<0.005
MMSE (ordinal)	1.93	1.459–2.575	0.68	1.6279	1.207–2.196	0.34
Blood loss >2 g/dL	2.13	1.259–3.575	0.214	1.921	1.217–2.396	0.117
Rectal temperature<36						
<40 min	1.821	0.466-7.115	0.3890	1.369	0.481-2.306	0.284
>40 min	5.06	3.895–6.575	<0.0001	3.2095	2.395–4.302	0.0322

Hazard ratio (HR) estimated from Cox proportional hazard regress-on model. Confidence interval (CI) of the estimated HR.

Conclusions:

- Age>76, marital status, alcohol use disorders, preoperative urinary catheterization were found to be preoperative predictors for onset of POD, while hypothermia time was associated with POD severity.
- Reducing the operative time, usage of preheated rinsing liquids in association with specific treatment may decrease POD incidence, morbidity and hospitalization costs.