Lichtheimia corymbifera: elucidating the mechanisms for iron acquisition and transporter during infection

Felicia Adelina Stanford^{1,2}, Volker U. Schwartz^{1,2} and Kerstin Voigt^{1,2}

¹ Leibniz Institute for Natural Product Research and Infection Biology, Hans-Knöll Institute (HKI), Jena Microbial Resource Collection (JMRC), Jena, Germany; ² Friedrich-Schiller-University Jena, Dept. Microbiology & Molecular Biology, Jena, Germany

I. Introduction

Pathogenic *Lichtheimia* species are clinically important, opportunistic fungi that can cause mucormycosis.^{1,2} Infections with mucoralean fungi are known for their rapid progression, diagnosing difficulties and the limited range of therapeutic options leading to high mortality rates.² Some of the major risk factors for the development of mucormycosis is elevated serum iron levels and the use of predisposing drugs (e.g. deferoxamine).^{2,3} Iron availability during infection is kept extremely low, with most of the iron bound to host proteins such as ferritin, transferrin and haemoglobin. This iron-withholding response in the host is a recognised example of nutritional-immunity and it is one of the main strategies used to restrict the growth of invading pathogens.^{3,4,5} As such, the specialised and highly efficient iron-uptake systems that fungi utilise in the environment to acquire iron i.e. the reductive pathway, siderophore uptake systems and sequestration of iron containing proteins (e.g. haem, hemin) are virulence factors that contribute to infection.^{5,6,7}

II. Objectives

To date, our knowledge regarding iron metabolism in *L. corymbifera* during infection remains unclear. As iron uptake is a key factor during infection, our main objectives are to determine the spectrum of iron sources *L. corymbifera* can utilise in the environment and identify which host iron sources are exploited. We also intend to identify which transcription factors are involved and gain insight into how these iron-uptake systems are regulated.

III. Results

Genomics-based identification of the Iron Acquisition Systems

	Pathway	Component	Copies
(A)	Reductive Iron Uptake	Iron Reductase	3
		Iron Permease (FTR1)	4
		Multicopper Oxidase	8
(B)	Haemoglobin transporter	Heme Oxygenase	2
(C)	Siderophores transporter	Deferoxamine Receptor (FOB1)	1
(D)	Iron Storage	Siderophore transporters	3
		NRPS-independent siderophore synthase	2
(E)	Regulation	Ferritin	2
		GATA transcription factor (SreA-like)	1

- Multiple copies of genes present in the iron uptake systems
- Broad range of functions may be possible

Iron Exploitation: Host Sources

- L. corymbifera* can utilise host ferritin and haemoglobin

Putative Siderophore Transporters and Siderophore utilisation

- Siderophore transporters show strong similarities to those of other fungi
- Utilisation of a broad spectrum of siderophores

Iron Regulation by GATA-type transcription factor

- Sequence similarities to GATA-type transcription regulators involved in iron adaptation in *A. fumigatus* and *H. capsulatum*
- Over four-fold up regulation under iron depletion

IV. Summary and Future Work

Our genomics, transcriptomics and physiological studies indicate that *L. corymbifera* is equipped with multiple copies of the genes involved in the three characterised iron uptake-systems. Our results show that *L. corymbifera* is able to utilise iron from numerous sources, including iron-containing host proteins. Our next aim is to identify which uptake systems play a key role in iron acquisition during infection. We intend to investigate the functionality of the multiple copies of iron uptake genes in the reductive pathway and the siderophore transporters. In addition, we also aim to determine the regulation of these genes. This will allow us to examine the significance of multiple gene copies in the iron uptake systems during infection.

Answering these key questions will aid our understanding of how these transporter systems contribute to virulence. Collectively, our findings have the potential to aid future research into the development of therapeutics.

References

- Ibrahim, A.S., (2014) Host-iron assimilation: pathogenesis and novel therapies of mucormycosis. *Mycoses*. 57(3):13-17
- Schwartz, V.U., Winter, S., Shelest, E., et al (2014) Gene Expansion Shapes Genome Architecture in the Human Pathogen *Lichtheimia corymbifera*: An Evolutionary Genomics Analysis in the Ancient Terrestrial Mucorales (Mucoromycotina). *PLoS Genetics* 10(8): e1004496. doi:10.1371/journal.pgen.1004496
- Almeida, R.S., Brunke, S., Albrecht, A., et al. (2008) The Hyphal-Associated Adhesin and Invasin Als3 of *Candida albicans* mediates iron acquisition from host ferritin. *PLoS Pathog* 4(11):e1000217. doi:10.1371/journal.ppat.1000217
- Ballou, E.R., and Wilson, D., (2016) The role of zinc and copper sensing in fungal pathogenesis. *Current Opinions in Microbiology*. 32:128-134
- Gerwien, F., Salyan, A., Wisgott, Brunke, S., Kasper, L., Hube, B. (2017) The Pathogen *Candida glabrata* does not depend on surface ferric reductases for iron acquisition. *Front. Microbiol.* 8:1055. doi:10.3389/fmicb.2017.01055
- Ibrahim, A.I., Gebremariam, T., Lin, L., Lou, G. et al. (2010) The high affinity iron permease is a key virulence factor required for *Rhizopus oryzae* pathogenesis. *Molecular Microbiology*. 77(3): 587-604
- Chen, L.H., Yang, S.L. and Chung, K.R., (2014) Resistance to oxidative stress via regulating siderophore-mediated iron acquisition by the citrus fungal pathogen *Alternaria alternata*. *Microbiology*. 160:970-979

Acknowledgments

This work is supported by the Jena School of Microbial Communications (JSMC), Friedrich-Schiller-University and funded by the German Academic Exchange (DAAD) – Deutscher Akademischer Austauschdienst and the German Research Association (CRC/TR124, ProjectA6)