ESMOLOL INCREASES NITRIC OXIDE AVAILABILITY AND E-NOS EXPRESSION IN A PORCINE MODEL OF LUNG RESECTION SURGERY

Ramos S¹, Puig A¹, Fernandez-Peña A¹, Simon C², Vara E³, Rancan L³
1 Gregorio Marañón General Hospital, Dept of Anaesthesiology & Intensive Care, Madrid, Spain,
2 Gregorio Marañón General Hospital, Dept of Thoracic Surgery, Madrid, Spain
3 Universidad Complutense de Madrid, Dept of Biochemistry and Molecular Biology, Madrid, Spain

Background and Goal of Study: lung resection and one lung ventilation (OLV) promote inflammatory response and secondary endothelial dysfunction partially due to a decrease in NO levels. Several beta blockers, as nebivolol, have shown to play a role in NO metabolism. However, there is little evidence regarding esmolol, the most widely used betablocker in the surgical setting. Our goal was to study the effects of esmolol on NO metabolism in a lung resection porcine model.

Materials and Methods:

24 minipigs → **Control (C) (n=8)**
Esmolol (E) (n=8)
Sham (S) (n=8)

Left thoracotomy	Lung Resection Surgery	Esmolol (0.5 mg/kg + 0.05 mg/kg/h)
✓	✓	
✓	✓	✓
✓

NO levels	Griess test
iNOS and eNOS protein expression	Western Blot
iNOS and eNOS mRNA levels	RT-PCR
Statistical analysis	ANOVA Mann-Whitney test

Results:

Conclusion: Our results suggest that esmolol could play an important role in NO metabolism downregulating eNOS and increasing NO availability. It may therefore prevent endothelial dysfunction associated with inflammatory responses by LRS and OLV.