ATTENTION-DEFICIT/HYPERACTIVITY DISORDER - - PREDICTORS OF PERSISTENCE INTO ADULTHOOD

Pereira, D.¹, Moreno, M.¹, Pereira, J.¹, Mendes, M.¹, Fernandes, R.¹, Ponte, A.¹, Viveiros, V.²
¹Psychiatry Resident, Centro Hospitalar Psiquiátrico de Lisboa;
²Psychiatrist, Centro Hospitalar Psiquiátrico de Lisboa

dmr_pereira@hotmail.com

INTRODUCTION

Attention deficit/hyperactivity disorder (ADHD) is a persistent neurodevelopmental condition characterized by persisting inattention, hyperactivity and/or impulsivity with significantly impaired functioning and development. Follow-up studies assessing the level of persistence of ADHD from childhood into adolescence and adulthood have been inconsistent, with ADHD prevalence between adults ranging from as low as 4% to as high as 76%^{1,2,7,8}. There are many studies documenting adverse outcomes from those affected by ADHD when compared to those without the disorder. ADHD affects a wide range of functional domains including academic, social and occupational contexts^{7,8}. Although these adverse adult outcomes associated with ADHD are unquestionable, much less clear are their childhood predictors.

Our objective is to identify the current knowledge in terms of factors associated with ADHD persistence or remission from childhood to adulthood through a systematic review of literature. A PubMed research were performed in January 2019 using the search keywords “ADHD”, “*predictors of persistence*”, “*risk factors*” and “*adulthood*”.

RESULTS

Several factors have been suggested as predictors of the course of ADHD. Most authors who evaluated the effect of socio-demographic features like gender, socioeconomic status at childhood and intelligence quotient were unanimous that these factors were not contributors to the persistence of ADHD through childhood to adulthood^{1,2,3,4,5}.

Four studies reported that persisters had higher levels of impairment^{1,2,5,6}. Also, individuals who were treated for ADHD in childhood were at a significantly higher risk for the persistence of the disease into adulthood. This effect remained significant even after controlling for ADHD severity^{4,5}.

Comorbid conditions like conduct disorder, major depressive disorder⁶, oppositional defiant disorder⁴ and anxiety disorders² have been suggested as influencing the outcome in ADHD subjects. However, only conduct disorder and major depressive disorder remained significant predictors after controlling for childhood ADHD symptoms.

The available literature indicates that ADHD severity, comorbid conduct disorder and major depressive disorder, and the treatment for ADHD are the main predictors of ADHD persistence from childhood to adulthood.

Other important issues, such as exposure to trauma and adversities, genetic factors and perinatal factors, have been reported as being significant predictors of persistence into adulthood. However, there is a limited number of studies evaluating this relationship and those available suffer from many limitations^{3,5,6}.

CONCLUSION

Even in light of the best available evidence, it is impossible for a clinician to determine whether a child presenting with ADHD will sustain the condition through adolescence and adulthood. The determination of this factors could help the delineation of prevention strategies in this group of patients. More prospective population-based cohort studies are important to expand our knowledge about ADHD trajectories across the life cycle.

REFERENCES

1. Russel A, Barkley MF (2007) ADHD in adults: what the science says. Guilford Press, New York
2. Biederman J, Petty CR, O'Connor KB, Hyder LL, Faraone SV (2011). Predictors of persistent ADHD: an 11-year follow-up study. J Psychiatr Res 45(2):150-155.
3. Biederman J, Petty CR, O'Connor KB, Hyder LL, Faraone SV (2012). Predictors of persistence in girls with attention deficit/hyperactivity disorder: results from an 11-year controlled follow-up study. Acta Psychiatr Scand 125(2):147-156.
4. Chang WLY, Qian Q, Tang H, Wang Y (2011). Related factors of early adulthood attention deficit hyperactivity disorder. Chin Ment Health J 25(12):5
5. Kessler RC, Adler LA, Barkley R, Biederman J, Conners CK, Faraone SV, Greenhill LL, Jaeger S, Secnik K, Spencer T, Ustun TB, Zaslavsky AM (2005). Patterns and predictors of attention-deficit/hyperactivity disorder persistence into adulthood: results from the national comorbidity survey replication. Biol Psychiatry 57(11):1442-1451.
6. Lara C, Fayyad J, de Graaf R, Kessler RC, Aguilar-Gaxiola S, Angermeyer M, Demyttenaere K, de Girolamo G, Haro JM, Jin R, Karam EG, Lepine JP, Mora ME, Ormel J, Posada-Villa J, Sampson N (2009) Childhood predictors of adult attention-deficit/hyperactivity disorder: results from the World Health Organization World Mental Health Survey Initiative. Biol Psychiatry 65(1):46-54.
7. Faraone SV, Asherson P, Banaschewski T, Biederman J, Buitelaar JK, Ramos-Quiroga JA, et al. Attention-deficit/hyperactivity disorder. Nat rRev Dis Primers. 2015;1:15020.
8. Asherson P, Buitelaar J, Faraone SV, Rohde LA. Adult attention-deficit hyperactivity disorder: key conceptual issues. Lancet Psychiatry. 2016;3(6):568-78.