Analgesic Requirements in Super Morbid and Morbid Obese Bariatric Patients Undergoing Laparoscopic Sleeve Gastrectomy: A Prospective Observational Study

Hamed ELGENDY, MD, PhD ^{1,2,3}; Soha ELMORSY, MD, PhD ^{2,5}; Talha YOUSSEF , MBBS ⁶ ; Ahmad BANJAR, MBBS ^{7,8}

1 Department of Anesthesia, Assiut University Hospitals, Egypt & 2 Anesthesia Dept., HAMAD Medical Corporation, Doha, Qatar & 3 Clinical Ass. Prof. Weill Cornell Medicine Qatar

2 Department of Medical Pharmacology Faculty of Medicine, Cairo University; 5 Research Consultant, King Abdullah Medical City Research Center, Makkah, Saudi Arabia;

6 Internal Medicine Dept., Prince Mohammad Bin Abdul-Aziz Hospital, Ministry of National Guard, Al Madinah, Saudi Arabia.

7Medical Student trainee , Umm Al Qura University & 8 King Abdullah Medical City; Saudi Arabia.

Background and aims: Opioid analgesia is associated with serious adverse effects; hypoventilation, hypoxaemia and ileus. Our aim was to investigate analgesic requirements for super morbid obese, (SMO), (BMI ≥ 50 kg/m2) when compared to morbid (MO), (BMI 40-49.9 kg/m2) individuals underwent sleeve gastrectomy and its impact on postoperative outcome.

Methods: We studied 297 consecutive patients (183 MO, 96 SMO) underwent bariatric surgery. Data analysis included perioperative anesthetic management.

Results: SMO patients exhibited higher asthmatics percentage (P = 0.016) and ASA III (P < 0.001). Lower serum albumin (P = 0.001) and higher INR levels (P = 0.020) were observed in SMO patients (Table 1).

Regular laryngoscope was commonly used in MO group, while other devices was higher in SMO, (P = 0.003). Higher propofol consumption, longer anesthesia time, increased crystalloid intake and urine output were observed in SMO group, respectively, (P = 0.001), (P = 0.033), (P < 0.001), (P = 0.017) (Table 2). Total morphine equivalent consumption dosages and VAS score were comparable in both groups at admission, stay or leaving PACU. Vomiting was higher in MO group, (P = 0.004). ICU admission was higher at SMO group (P < 0.001), however, PACU stay time was comparable, (P =0.060) (Table 3).

Conclusions: We found no differences in perioperative analgesics consumption or postoperative outcome in between the studied groups except longer ICU stay, prolonged anesthesia time, increased crystalloid intake and urine output.

Table -1: Demographic Characteristic

Characteristics	Morbidly Obese (n=183)	Super- Morbidly Obese (n=96)	P-Value
Age/ Years	34.5 ± 10.4	34.1 ± 10.7	0.753
Gender n (%)			
Male	72/183 (39.3)	46/96 (47.9)	0.169
Female	111/183 (60.7)	50/96 (52.1)	0.169
Weight/ Kg	118.8 ± 18.0	151.6 ± 23.1	< 0.001
Height/m	1.7 ± 0.1	1.6 ± 0.1	0.005
Malamprat Classification n (%)			
I			
II	40/183 (21.9)	24/96 (25.0)	0.835
III	114/183 (62.3)	57/96 (59.4)	
	29/183 (15.8)	15/96 (15.6)	
ASA Classification n (%)			
Class II	39/183 (21.3)	0/96 (0.0)	< 0.001
Class III	144/183 (78.7)	96/96 (100.0)	< 0.001
Preoperative Laboratory parameters			
Total Bilirubin mg/dl	0.4 ± 0.3	0.5 ± 0.3	0.208
ALT (IU/L)	46.2 ± 13.2	47.6 ± 25.5	0.928
AST (IU/L)	24.9 ± 13.2	26.3 ± 18.8	0.694
Magnesium (mg/dl)	2.0 ± 0.5	1.9 ± 0.5	0.738
Calcium (mg/dl)	8.8 ± 1.2	8.8 ± 1.2	0.143
Sodium (mmol/l)	136.8 ± 3.3	136.7 ± 2.2	0.699
Potassium (mmol/l)	4.3 ± 0.3	4.6 ± 1.1	0.247
BUN (mg/dl)	13.9 ± 15.4	13.9 ± 16.0	0.806
Creatinine (mg/dl)	0.8 ± 0.2	0.8 ± 0.2	0.073
Albumin (g/dl)	3.7 ± 0.6	3.6 ± 0.8	0.001
Platelet count (× 10 ⁹ /L)	288.9 ± 67.7	301.8 ± 80.5	0.915
INR	1.00 ± 0.09	1.01 ± 0.12	0.020
PT (Sec)	12.6 ± 1.6	12.5 ± 1.2	0.723
APTT (Sec)	32.8 ± 6.6	33.4 ± 7.3	0.724
Hemoglobin (g/dl)	13.6 ± 1.9	13.4 ± 1.7	0.665
WBC count (× 10 ⁹ /L)	8.4 ± 2.0	8.9 ± 2.3	0.162
Co- Morbidities n (%)			
HTN	36/183 (19.7)	20/96 (20.8)	0.818
DM	34/183 (18.6)	19/96 (19.8)	0.806
Renal	5/183 (2.7)	0/96 (0.0)	0.102
CNS (Epilepsy)	0/183 (0.0)	3/96 (3.1)	0.016
CVS	6/183 (3.3)	7/96 (7.3)	0.131
Respiratory (Bronchial asthma)	7/183 (3.8)	11/96 (11.5)	0.014
Associated Therapy n (%)			
Anti HTN	31/183 (16.9)	18/96 (18.8)	0.706
Insulin	09/183 (4.9)	05/96 (5.2)	0.916
Oral hypoglycemic	28/183 (15.8)	16/96 (16.7)	0.860
Anti coagulant drugs	08/183 (4.4)	02/96 (2.1)	0.329
Anti Asthmatic drugs	03/183 (1.6)	03/96 (3.1)	0.416
Smoking	40/183 (21.9)	25/96 (26.0)	0.432
Alcohol	1/183 (0.5)	0/96 (0.0)	0.468

Values are either mean ± SD or number and percentage. P < 0.05 considered statistically significant.

Table -2: Intra Operative Variables

Characteristics	Morbidly Obese (n=183)	Super-Morbidly Obese (n=96)	P- Value
Premedication : Midazolam / mg	0.3 ± 0.9	0.2 ± 0.6	0.508
IV anaesthetics: Propofol / mg	212.6 ± 59.4	238.1 ± 64.0	0.001
Intra Operative analgesics Usage_ n (%)			
Paracetamol	148 /183 (80.9)	75/96 (78.1)	0.586
Morphine	146 /183 (79.8)	77/96 (80.2)	0.933
Intra Operative agents Usage_ n (%)			
Succinylcholine	20 /183 (10.9)	11/96 (11.5)	0.894
Atracurium	98 /183 (53.6)	47/96 (49.0)	0.466
Rocuronium	87 /183 (47.5)	51/96 (53.1)	0.375
Neostigmine	123 /183 (67.2)	58/96 (60.4)	0.259
Glycopyrrolate	109 /183 (59.6)	50/96 (52.1)	0.231
Atropine	31 /183 (16.9)	11/96 (11.5)	0.224
Sevoflurane	186 /183 (91.8)	89/96 (92.7)	0.790
Desflurane	15 /183 (8.3)	07/96 (7.3)	0.650
Ranitidine	109 /183 (59.6)	49/96 (51.0)	0.172
Dexamethasone	120/183 (65.6)	68/96 (70.8)	0.373
Ondansetron	144 /183 (78.7)	78/96 (81.3)	0.614
Intra Operative techniques n (%)			
Difficult Intubation	08/183 (4.4)	08/96 (8.3)	0.176
Cormack- Lehane grade			
Grade I	93/183 (50.8)	44/96 (45.8)	0.639
Grade II	74/183 (40.4)	46/96 (47.9)	
Grade III	14/183 (7.7)	05/96 (5.2)	
Grade IV	02/183 (1.1)	01/96 (1.0)	
Airway intubation Method n (%)			
- Regular laryngoscope	138/183 (75.4)	57/96 (59.4)	0.033
- Video - assisted	37/183 (20.2)	29/96 (30.2)	
- fibre-optic Intubation	05/183 (2.7)	06/96 (6.3)	
-Retro molar device	03/183 (1.6)	04/96 (4.2)	
Crystalloids / ml	2293.7 ± 778.1	2694.8 ± 835.8	< 0.001
Blood Loss / ml	19.8 ± 7.4	59.0 ± 21.2	0.367
Urine Output / ml	103.4 ± 72.9	190.1 ± 85.2	0.017
Anesthesia time/min	127.7 ± 38.1	137.6 ± 42.6	0.033
Surgery time/min	102.7 ± 35.1	107.2 ± 39.3	0.366

Values are either mean ± SD or number. and percentage. P < 0.05 considered statistically significant.

Table- 3: Post operative Variables

Characteristics	Morbidly Obese (n=183)	Super- Morbidly Obese (n=96)	P value
PACU VAS Score			
Admission	2.59 ± 1.62	2.43 ± 1.53	0.558
Worst	3.19 ± 1.80	3.21 ± 1.86	0.959
Leaving	1.86 ± 1.07	1.70 ± 1.02	0.176
PACU analgesic usage n (%)			
Morphine	34 /183 (18.6)	19/96 (19.6)	0.806
Fentanyl	24 /183 (13.1)	14/96 (14.6)	0.734
Tramadol	42 /183 (23.0)	20/96 (20.8)	0.686
Paracetamol	41 /183 (22.4)	21/96 (21.9)	0.920
Lornoxicam	15 /183 (2.8)	9/96 (9.4)	0.739
ketorolac	1 /183 (0.5)	0/96 (0.0)	0.468
Total Morphine equivalent	10.5 ± 4.5	10.6 ± 4.4	0.878
PONV & treatment n (%)			
Nausea	29 /183 (15.8)	13/96 (13.5)	0.609
Vomiting	18 /183 (9.8)	3/96 (3.1)	0.044
IV Ondansetron	23 /183 (12.6)	12/96 (12.5)	0.987
IV Metoclopramide	38 /183 (20.8)	14/96 (14.6)	0.208
PACU stay time/ min.	90.4 ± 41.0	97.3 ± 31.9	0.060
ICU admission n (%)	03 /183 (1.6)	12/96 (12.5)	< 0.001

Values are either mean ± SD or number and percentage. P < 0.05 considered statistically significant.