QUALITY OF LIFE AND ORAL PAIN IN HEAD AND NECK CANCER PATIENTS RECEIVING RADIO/CHEMORADIO THERAPY

Maria Kouri¹, Ourania Nicolatou-Galitis¹, Athina Vadalouca², Vasilios Kouloulis³, Georgios Maropoulos⁴, Panagiotis Kanellopoulos⁴, Efthymios Kyrodimos⁵, Erofilis Papadopoulou¹, Emmanouil Vardas¹, Evangelos Galitis⁶ Miltiadis Trichas⁷, Anastasia Vourli¹, Anna Zigogianni⁸, Zoi Liakouli⁸, Amanda Psyrris⁹, Ioannis Kouvaris⁸

¹Clinic of Hospital Dentistry, School of Dentistry, National and Kapodistrian University of Athens, Athens, Greece
²Pain Clinic, “Athens Medical Center”, Athens, Greece
³2nd Radiology Department, Radiotherapy Unit, Medical School, National and Kapodistrian University of Athens, Athens Greece
⁴Department of Biochemistry, “Laiko” General Hospital of Athens, Athens, Greece
⁵ENT Clinic “Hippokrateio” Hospital, Medical School, National and Kapodistrian University of Athens, Athens Greece
⁶Clinic of Orofacial Pain, School of Dentistry, National and Kapodistrian University of Athens, Athens, Greece
⁷Radiotherapy Department, “Iaso” Hospital, Athens, Greece
⁸1st Radiology Department, Radiotherapy Unit, Medical School, “Areteiaion” University Hospital, National and Kapodistrian University of Athens, Greece
⁹Oncology Department, “Attikon” University Hospital, Medical School, National and Kapodistrian University of Athens, Athens Greece

OBJECTIVES:

To assess pain related changes on Quality of Life (QoL), in Head/Neck cancer patients receiving radio/chemoradiotherapy.

PATIENTS & METHODS:

QoL ASSESSMENT:

Core-30 (EORTC QLQ-C30)V3.0
Head and Neck module (QLQ-H&N35)
QoL was Assessed:
before radiotherapy (t0), 5th-7th week (t1).

ORAL PAIN EVALUATION:

Verbal Analogue Scale (VAS).
Scores: 0 – 10;
1 - 4: mild, 5 – 7: moderate,
8 – 10: severe.

PARAMETERS		N=30	%
Gender	Male	26	86.7
	Female	4	13.3
Age (Mean)		64.3	
Radiotherapy	Postoperative	17	56.7
	Radical	13	43.3
Total Dose (Mean, Gy)		65.38	
Concomitant Chemotherapy		23	76.7
Tumor Site	Oral/ Oropharyngeal	14	46.7
	Laryngeal	6	20
	Parotid	3	10
	Other	7	23.3

CONCLUSIONS:

Pain was related to most of the QoL score changes. Appropriate interventions can be directed towards managing the pain and improving QoL of the patients.

RESULTS:

PARAMETERS			
PAIN	N	QoL Values Core-30 (EORTC QLQ-C30)	Paired t-test
Severe	7	Global QoL/ Physical/ Role / Social Functions	<0.006
		Fatigue/ Dyspnoea/ Appetite Loss	<0.004
Moderate	10	Physical/ Role/ Emotional/ Cognitive/ Social Functions	<0.017
		Fatigue/ Dyspnoea/ Appetite Loss/ Constipation	<0.029
Mild	9	Physical/ Role / Social Functions	<0.038
		Fatigue/ Appetite Loss	<0.039

PARAMETERS			
PAIN	N	QoL Values Head and Neck module (QLQ-H&N35)	Paired t-test
Severe	7	Swallowing/ Dry Mouth/ Opening Mouth	0.000
		Senses/ Social Eating/ Sticky Saliva	0.001
		Speech/ Social Contact/ Sexuality/ Felt Ill/ Weight Loss	<0.030
Moderate	10	Swallowing/ Senses/ Social Eating/ Dry Mouth/ Sticky Saliva	<0.007
		Social Contact/ Sexuality/ Felt Ill/ Weight Loss	<0.029
Mild	9	Speech/ Social Eating/Sticky Saliva/ Weight Loss	<0.004
		Senses/ Dry mouth/ Social Contact	<0.035

Severe pain: Global QoL, Physical, Role, Social Functions were significantly decreased (p<0.006) (t1 vs t0 assessment). Insignificant reduction of Emotional and Cognitive Functions; Symptom Scales and H&N35 scores were significantly increased. Nausea/Vomiting, Insomnia, Constipation, Financial problems, Diarrhoea, Teeth problems, Coughing and Weight gain, statistically insignificant.
Moderate pain: Significantly lower Functional Scales scores (p<0.02) (t1 vs t0 assessment). Insignificant reduction of Global QoL was (p=0.160); Symptom Scales and H&N35 scores were significantly increased, except for Nausea/Vomiting, Insomnia, Financial problems, Diarrhoea, Speech and Teeth problems, Opening mouth, Coughing and Weight gain (statistically insignificant).